

08-14-024 Rec'd PCT/PTO 10 AUG 2000


U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER 256/137 US
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. APPLICATION NO. (If known, see 37 CFR 1.5) Not Assigned 09/622101
INTERNATIONAL APPLICATION NO. PCT/EP99/00945	INTERNATIONAL FILING DATE 12 February 1999	PRIORITY DATE CLAIMED 13 February 1998
TITLE OF INVENTION NOVEL MEANS AND METHODS FOR THE PREPARATION AND ACTIVATION OF NUCLEOSIDE AND NUCLEOTIDE BASED DRUGS		
APPLICANT(S) FOR DO/EO/US Max-Planck-Gesellschaft zur Forderung der Wissenschaften e.V.		
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:		
<p>1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.</p> <p>2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.</p> <p>3. <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).</p> <p>4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.</p> <p>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))</p> <p>a. <input checked="" type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau).</p> <p>b. <input type="checkbox"/> has been transmitted by the International Bureau.</p> <p>c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</p> <p>6. <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)).</p> <p>7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))</p> <p>a. <input checked="" type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau).</p> <p>b. <input type="checkbox"/> have been transmitted by the International Bureau.</p> <p>c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</p> <p>d. <input type="checkbox"/> have not been made and will not be made.</p> <p>8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).</p> <p>9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).</p> <p>10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</p>		
Items 11. to 16. below concern document(s) or information included:		
<p>11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</p> <p>12. <input type="checkbox"/> An assignment document for recording.</p> <p>13. <input type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment.</p> <p>14. <input type="checkbox"/> A substitute specification.</p> <p>15. <input type="checkbox"/> A change of power of attorney and/or address letter.</p> <p>16. <input checked="" type="checkbox"/> Other items or information:</p> <p>PCT Request</p> <p>PCT Written Opinion</p> <p>PCT Cover page of International Application Published Under the Patent Cooperation Treaty (PCT)</p> <p>PCT Notification of Transmittal of the International Search Report or the Declaration</p> <p>PCT Notification of Transmittal of the International Preliminary Examination Report</p> <p>Return Postcard</p> <p>Check for \$970.00</p>		

Express Mail mailing label number EL 712165831 US

Date of Deposit 8/10/00

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Commissioner of Patents and Trade marks, Washington, DC 20231.

Amanda Halverson
Type or Printed Name of Person
Mailing Paper or Fee
Amanda Halverson
Signature

U.S. APPLICATION NO. (If known, fee 37 CFR 1.51) Not 299 Assigned 2101 INTERNATIONAL APPLICATION NO. PCT/EP99/00945		ATTORNEY'S DOCKET NUMBER 256/137 US	
17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) : Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$970.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$840.00 International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$690.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$670.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(1)-(4) \$96.00 ENTER APPROPRIATE BASIC FEE AMOUNT =		CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).		\$ 840.00 \$ 130.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	- 20 =		X \$18.00
Independent claims	- 3 =		X \$78.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$260.00
TOTAL OF ABOVE CALCULATIONS =			\$ 970.00
Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).			\$
SUBTOTAL =			\$
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).			\$
TOTAL NATIONAL FEE =			\$
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property			+
TOTAL FEES ENCLOSED =			\$ 970.00
			Amount to be refunded:
			charged:
a. <input checked="" type="checkbox"/> A check in the amount of \$ <u>970.00</u> to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>12-2475</u> . A duplicate copy of this sheet is enclosed.			
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.			
SEND ALL CORRESPONDENCE TO: Suzanne L. Biggs Lyon & Lyon 633 West Fifth Street, Suite 4700 Los Angeles, CA 90071			
			SIGNATURE:  Suzanne L. Biggs
			NAME 30,158
			REGISTRATION NUMBER August 10/2000

Rec'd PCT/PTO 29 JAN 2001

Patent

Attorney Docket: 256/137

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of:

Roger S. Gooding, Manfred
Konrad, Arnon Lavie, Joachim
Reinstein, and Ilme Schlichting

Serial No.: 09/622,101

Filed: 08/10/00

I.A. Filing Date: 02/12/99

Priority Date: 02/13/98

For: NOVEL MEANS AND METHODS
FOR THE PREPARATION AND
ACTIVATION OF NUCLEOSIDE
AND NUCLEOTIDE BASED DRUGS

Group Art Unit: Not yet
assigned

Examiner: Not yet assigned

PRELIMINARY AMENDMENT

BOX MISSING PARTS
Commissioner for Patents
Washington, D.C. 20231

Sir:

Please enter the following amendments before examining this
application.

SD-159287.1

CERTIFICATE OF MAILING
(37 C.F.R. §1.10(a))

I hereby certify that this paper (along with any referred to as being attached or enclosed) is being deposited with the United States Postal Service on the date shown below with sufficient postage as 'Express Mail Post Office to Addressee' in an envelope addressed to the Commissioner for Patents, Washington, D.C. 20231.

EL356078344US
Express Mail Label No.

January 29, 2001
Date of Deposit

Vanessa Hinton
Name of Person Mailing Paper

Vanessa Hinton
Signature of Person Mailing Paper

In the Claims:

Please amend claims 3, 5, 7 to 14, 17, 18, 20, 21, 23, 25 to 28, 32, 34, and 36 as shown on the attached "Claim Sheets Marked up to Show Changes".

Please add new claims 39 to 42 as shown in the attached Clean Claim Sheets which show all claims pending in the application, after entry of this amendment.

REMARKS

This preliminary amendment is being submitted concurrently with Applicants' Response to the Notice to File Missing Parts. Applicants submit that this preliminary amendment is appropriate under 37 C.F.R. § 1.115 and request that it be entered.

This application us the United States national phase application corresponding to PCT/EP99/00945.

Claims 3, 5, 7 to 14, 17, 18, 20, 21, 23, 25 to 28, 32, 34 and 36 have been amended to eliminate certain multiple dependencies and references to multiple dependencies in multiple dependent claims and to put the claims in better form for examination under United States practice. Applicants submit that these amendments are supported by the specification and give rise to no issue of new matter.

New claims 39 to 42 have been added to more particularly point out and distinctively claim certain aspects of Applicants' invention. Applicants submit that these claims are clearly supported by the specification and give rise to no issue of new matter.


If the Examiner believes that a telephone interview would expedite prosecution of this application, he is encouraged to telephone the undersigned Applicants' attorney.

If the fees submitted are incorrect or if any other fees are due in connection with this submission, please charge any fee or credit any overpayment to Deposit Account No. 12-2475.

Respectfully submitted,

LYON & LYON LLP

Dated: January 29, 2001

By: 
Suzanne L. Biggs
Reg. No. 30,158


633 West Fifth Street, Suite 4700
Los Angeles, California 90071-2066
(213) 489-1600

Serial No. 09/622,101

Clean Claim Sheets

1. A method for the production of a polypeptide having or having enhanced kinase activity for a nucleoside or nucleotide analog, said method comprising substituting, adding or deleting at least one amino acid of a protein having nucleoside or nucleotide kinase activity at a position in the protein where:
 - (a) the amino acid is at position X_2 and/or X_3 in the consensus sequence $GX_1X_2X_3X_4GK$ of the P-loop;
 - (b) the amino acid is in the LID region; and/or
 - (c) the amino acid is at position 105 in the amino acid sequence of human thymidylate kinase or at a corresponding position in a protein having nucleoside or nucleotide kinase activity.
2. The method of claim 1, wherein said nucleoside is adenosine, cytidine, guanosine, thymidine or uridine or based on any of these.
3. (Amended) The method of claim 1, wherein said nucleotide is a nucleoside monophosphate.
4. The method of claim 3, wherein said nucleoside monophosphate is thymidylate.
5. (Amended) The method of claim 3, wherein said protein is derived from a eukaryotic or prokaryotic organism.
6. The method of claim 5, wherein said organism is human or a yeast.

7. (Amended) The method of claim 5, wherein said protein comprises the amino acid sequence of any one of SEQ ID NOS: 1 to 13 or a fragment thereof.
8. (Amended) The method of claim 5, wherein said amino acid which is substituted or added in (a) is glutamic acid, glycine or lysine and the amino acid substituted in (c) is tyrosine.
9. (Amended) The method of claim 8, wherein said amino acid which is substituted or added in (b) is a basic amino acid, preferably arginine.
10. (Amended) The method of claim 9, where the LID has the consensus sequence R/KXXXXXERYEXXXQ.
11. (Amended) The method of claim 10, wherein said polypeptide exhibits kinase activity for a nucleoside or nucleotide analog which is higher than that of a corresponding wild type eukaryotic enzyme.
12. (Amended) The method of claim 10, wherein said nucleoside analog is AZT, d4T or has the following structure:



wherein B is any nucleobase or analog thereof, and X is O, CH₂, NH or S.

13. (Amended) The method of claim 12, wherein the amino acid substitution(s) result in the P-loop and/or in the LID region of a bacterial nucleoside or nucleotide kinase.
14. (Amended) A polynucleotide encoding the polypeptide obtainable by the method of claim 10.
15. A vector containing the polynucleotide of claim 14.
16. The vector of claim 15, wherein the polynucleotide is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic cells.
17. (Amended) The vector of claim 16, which is a gene transfer or a gene targeting vector.
18. (Amended) A host cell genetically engineered with the vector of claim 16.
19. A method for producing a polypeptide having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog comprising
 - (a) culturing the host cell of claim 18, and
 - (b) recovering said polypeptide from the culture.
20. (Amended) A method for producing cells capable of expressing a polypeptide having nucleoside or nucleotide kinase activity for nucleoside or nucleotide analogs

comprising genetically engineering cells with the vector of claim 16.

21. (Amended) A polypeptide having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog encoded by a polynucleotide of claim 14 or a biologically active fragment thereof.
22. An antibody specifically recognizing the polypeptide of claim 21.
23. (Amended) A composition comprising
 - (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide; or
 - (b) the polypeptide of claim 21;
 - (c) optionally a nucleoside or nucleotide analog; and
 - (d) optionally a pharmaceutically acceptable carrier.
24. The composition of claim 23, wherein said protein is a bacterial nucleoside or nucleotide kinase.
25. (Amended) The composition of claim 24, wherein said protein is a bacterial TmpK.
26. (Amended) The composition claim 25, wherein said protein has at least the P-loop and/or the LID region of E.coli TmpK.

27. (Amended) The composition of claim 25, wherein said TmpK comprises the amino acid sequence shown in SEQ ID NO: 4 or a biologically active fragment thereof.
28. (Amended) A kit comprising the vector of claim 16 and optionally a nucleoside or nucleotide analog.
29. A method for identifying an inhibitor of a nucleoside or nucleotide kinase comprising the steps of:
- (a) contacting the polypeptide of claim 21 or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a compound to be screened under conditions that permit binding of said compound to the nucleoside or nucleotide kinase, and
 - (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the absence or decrease of the signal is indicative for an inhibitor of a nucleoside or nucleotide kinase.
30. A method for identifying a nucleoside or nucleotide based prodrug comprising the steps of
- (a) contacting the polypeptide of claim 21 or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a nucleoside or nucleotide analog to be screened under conditions that permit kinase activity of said polypeptide, and
 - (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the presence of a signal is indicative for a putative prodrug.

31. A method for the production of a pharmaceutical composition comprising the steps of

- (a) contacting the polypeptide of claim 21 or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a compound to be screened under conditions that permit binding of said compound to the nucleoside or nucleotide kinase, and
- (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the absence or decrease of the signal is indicative for an inhibitor of a nucleoside or nucleotide kinase, or
- (a') contacting the polypeptide of claim 21 or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a nucleoside or nucleotide analog to be screened under conditions that permit kinase activity of said polypeptide, and
- (b') detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the presence of a signal is indicative for a putative prodrug; and
- (c) formulating the inhibitor identified in step (b) or the nucleoside or nucleotide analog identified in step (b') in a pharmaceutically acceptable form.

32. (Amended) Use of

- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing

said protein *in vivo* or a vector containing said polynucleotide, or

(b) the polypeptide of claim 21

for the preparation of a pharmaceutical composition for the activation of nucleoside or nucleotide analogs or nucleoside or nucleotide based prodrugs.

33. The use of claim 32, wherein said activation results in a cytotoxic nucleoside or nucleotide.
34. (Amended) The use of claim 33, wherein said viral infection is HIV infection.
35. Use of the inhibitor obtainable by the method of claim 31 for the preparation of a pharmaceutical composition for inhibiting virus replication or for treating cancer.
36. (Amended) A method for the preparative synthesis of a nucleoside phosphate analog comprising:
- (a) using a polynucleotide of claim 14 in a noncellular system or in a cell *ex vivo*, and
 - (b) formulating the cells modified in step (a) in a pharmaceutically acceptable form.
37. The composition of any one of claims 23 to 27 or the use of any one of claims 32 to 34, wherein said composition is a pharmaceutical composition and further comprises or is designed to be administered with a nucleoside or nucleotide analog, preferably AZT or d4T.

38. Use of

- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide,
- (b) the polypeptide of claim 21, the polynucleotide of claim 14 or the vector of any one of claims 15 to 17 for the preparation of nucleoside phosphates or analogs and derivatives thereof.

39. The method of claim 10 wherein said protein comprises the amino acid sequence of any one of SEQ. ID. NOS. 1 to 13 or a fragment thereof.

40. The method of claim 11 wherein said enzyme is a human enzyme.

41. The method of claim 13 wherein said bacterial kinase is a Tmpk of E.coli.

42. The use according to claim 32 wherein said pharmaceutical composition is used for the treatment of a viral infection or disease for the treatment of cancer.

Serial No. 09/622,101

Claim Sheets Marked up to Show Changes

3. (Amended) The method of claim 1 [or 2], wherein said nucleotide is a nucleoside monophosphate.
5. (Amended) The method of [any one of claims 1 to 4] claim 3, wherein said protein is derived from a eukaryotic or prokaryotic organism.
7. (Amended) The method of [any one of claims 1 to 6] claim 5, wherein said protein comprises the amino acid sequence of any one of SEQ ID NOS: 1 to 13 or a fragment thereof.
8. (Amended) The method of [any one of claims 1 to 7] claim 5, wherein said amino acid which is substituted or added in (a) is glutamic acid, glycine or lysine and the amino acid substituted in (c) is tyrosine.
9. (Amended) The method of [any one of claims 1 to] claim 8, wherein said amino acid which is substituted or added in (b) is a basic amino acid, preferably arginine.
10. (Amended) The method of [any one of claims 1 to] claim 9, where the LID has the consensus sequence R/KXXXXXXERYEXXXQ.
11. (Amended) The method of [any one of claims 1 to] claim 10, wherein said polypeptide exhibits kinase activity for a nucleoside or nucleotide analog which is higher than that of [the/]a corresponding wild type eukaryotic[, preferably human] enzyme.

12. (Amended) The method of [any one of claims 1 to 11] claim 10, wherein said nucleoside analog is AZT, d4T or has the following structure:



wherein B is any nucleobase or analog thereof, and X is O, CH₂, NH or S.

13. (Amended) The method of [any one of claims 1 to] claim 12, wherein the amino acid substitution(s) result in the P-loop and/or in the LID region of a bacterial nucleoside or nucleotide kinase[, preferably those of the TmpK of E.coli].
14. (Amended) A polynucleotide encoding the polypeptide obtainable by the method of [any one of claims 1 to 13] claim 10.
17. (Amended) The vector of claim [15 or] 16, which is a gene transfer or a gene targeting vector.
18. (Amended) A host cell genetically engineered with the vector of [any one of claims 15 to 17] claim 16.
20. (Amended) A method for producing cells capable of expressing a polypeptide having nucleoside or nucleotide kinase activity for nucleoside or nucleotide analogs comprising genetically engineering cells with [the

polynucleotide of claim 14, or with] the vector of [any one of claims 15 to 17] claim 16.

21. (Amended) A polypeptide having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog encoded by a polynucleotide of claim 14[, obtainable by the method of any one of claims 1 to 13 or 19 or from cells produced by the method of claim 20] or [comprising] a biologically active fragment [of any of these] thereof.

23. (Amended) A composition comprising

- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide; or
- (b) the polypeptide of claim 21[, the polynucleotide of claim 14 or the vector of any one of claims 15 to 17];
- (c) optionally a nucleoside or nucleotide analog; and
- (d) optionally a pharmaceutically acceptable carrier.

25. (Amended) The composition of claim [23 or] 24, wherein said protein is a bacterial TmpK.

26. (Amended) The composition of [any one of claims 23 to] claim 25, wherein said protein has at least the P-loop and/or the LID region of E.coli TmpK.

27. (Amended) The composition of claim 25 [or 26], wherein said TmpK comprises the amino acid sequence shown in SEQ ID NO: 4 or a biologically active fragment thereof.

28. (Amended) A kit comprising [the polynucleotide of claim 14,] the vector of [any one of claims 15 to 17, the protein of claim 21 or the antibody of claim 22,] claim 16 and optionally a nucleoside or nucleotide analog.

32. (Amended) Use of

(a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide, or

(b) the polypeptide of claim 21[, the polynucleotide of claim 14 or the vector of any one of claims 15 to 17; and/or

(c) the nucleoside or nucleotide analog identified in the method of claim 30]

for the preparation of a pharmaceutical composition for the activation of nucleoside or nucleotide analogs or nucleoside or nucleotide based prodrugs [and/or for the treatment of viral infections and/or diseases or cancer].

34. (Amended) The use of claim [32 or] 33, wherein said viral infection is HIV infection.

36. (Amended) A method for the preparative synthesis of a nucleoside phosphate analog comprising:

(a) using a polynucleotide of claim 14 [or as defined in any one of claim 23 to 27] in a noncellular system or in a cell *ex vivo*, and

(b) formulating the cells modified in step (a) in a pharmaceutically acceptable form.

8/PRJS

Our Ref.: B 3270 PCT

**Novel means and methods for the preparation and activation of nucleoside
and nucleotide based drugs**

The present invention relates to novel means and methods for the preparation and activation of nucleoside and nucleotide based drugs. In particular, the present invention involves a method for the production of a polypeptide having or having enhanced kinase activity for a nucleoside or nucleotide analog and to polypeptides obtainable by said method. The present invention also relates to polynucleotides and vectors encoding said polypeptide obtainable by the method of the invention as well as to host cells transformed therewith. Antibodies against said polypeptide are also within the scope of the present invention. The present invention additionally relates to pharmaceutical and diagnostic compositions as well as kits comprising proteins having kinase activity for a nucleoside or nucleotide analog or the before described polypeptides, polynucleotides, vectors and antibodies. Furthermore, the present invention relates to the use of the before described proteins, polypeptides, polynucleotides, vectors and antibodies for the preparation of pharmaceutical compositions for treating, preventing and/or delaying a disease related to viral infection or cancer. In addition, the present invention relates to a method for identifying inhibitors of nucleoside or nucleotide kinases and to methods for identifying nucleoside or nucleotide based prodrugs employing the above mentioned polypeptides, polynucleotides, vectors and host cells. Also the invention relates to the compounds identifiable by said methods as well as to pharmaceutical and diagnostic compositions comprising said inhibitors. Moreover, the present invention relates to the use of proteins and polypeptides having nucleoside or nucleotide kinase activity or their encoding polynucleotides or vectors for the preparation of nucleoside or nucleotide phosphates or analogs and derivatives thereof.

Several documents are cited throughout the text of this specification. Each of the documents cited herein (including any manufacturer's specifications, instructions, etc.) are hereby incorporated herein by reference; however, there is no admission that any document cited is indeed prior art as to the present invention.

The class of medicinal agents termed prodrugs are compounds that exert a desired biological effect only after some sort of modification inside the body. AZT and d4T are examples of nucleoside analog prodrugs that attain their medicinal activity only after being phosphorylated three times to their triphosphate form by cellular kinases. Thus, the efficacy of these compounds is determined in part by the efficiency of the kinases that activate them. Compounds that are poorly activated (as is the case for AZT) do not achieve their full therapeutic potential. Moreover, the intermediate metabolites between the administered prodrug and the active triphosphate form can be toxic. Therefore, it is very important to use prodrugs that are efficiently transformed to their active form or to improve the activity of the rate limiting enzyme(s) in the activation pathway of a prodrug. The latter approach, however, was challenged in the art as being naive and doubtful to work at all; see, e.g., Balzarini, *Nature Medicine* 4 (1998), 2. Recently, Guettari, (*Virology* 235 (1997), 398-405) described the improvement of AZT metabolism by use of the Herpes Simplex virus-1 thymidilate kinase (HSV-1 TK) and suggested that gene transfer might be envisioned for genetic pharmacomodulation of anti-viral drugs. However, the extent to which HSV-1 TK improved AZT metabolism was only about 7-fold and no general method had been presented how to improve or create new kinase proteins or prodrugs that are suitable for a therapeutic approach. In two other recent reports (Lavie, *Nature Struct. Biol.* 4 (1997), 601—604; Lavie, *Nature Med.* 3 (1997), 922—924) it could be shown that the P-loop of the yeast TmpK is involved in limiting the conversion of AZT to AZTTP. The conclusion of crystallographic studies presented in these reports with the yeast TmpK enzyme have, however, been questioned, in particular whether other TmpK enzymes such as the human counterpart have the problem with the P-loop

observed with the yeast enzyme (Kenyon, Nature Struct. Biol. 4 (1997), 595—597). Hence, each enzyme seems to present a different situation, and no general method has been presented for modeling the catalytic domain of a kinase enzyme so as to obtain a high enzymatic activity with nucleoside or nucleotide analogs, such as AZT; nor was it even clear to what extent such method was required or possible.

Thus, the technical problem of the present invention is to provide means and methods for the preparation and activation of nucleoside and nucleotide based drugs.

The solution to this technical problem is achieved by providing the embodiments characterized in the claims.

Accordingly, the invention relates to a method for the production of a polypeptide having or having enhanced kinase activity for a nucleoside or nucleotide analog, said method comprising substituting, adding or deleting at least one amino acid of a protein having nucleoside or nucleotide kinase activity at a position in the protein where:

- (a) the amino acid is at position X_2 and/or X_3 in the consensus sequence $GX_1X_2X_3X_4GK$ of the P-loop;
- (b) the amino acid is in the LID region; and/or
- (c) the amino acid is at position 105 in the amino acid sequence of human thymidylate kinase or at a corresponding position in a protein having nucleoside or nucleotide kinase activity.

In context with the present invention, the term "nucleoside or nucleotide analog" refers to naturally occurring nucleosides that are modified at the sugar/base while nucleotide analogs may additionally have modifications, e.g., of the α -phosphate. Usually, said nucleoside is adenosine, cytidine, guanosine, thymidine or uridine or based on any of these, such as inosine. Nucleoside and nucleotide analogs useful in accordance with the present invention include 2', 3'-dideoxynucleoside analogs

and other analogs lacking a hydroxyl group equivalent to the 3'-OH of natural nucleoside in terms of their reactivity with reverse transcriptase or other DNA polymerases. Preferably said nucleotide is a nucleoside monophosphate, most preferably said nucleoside monophosphate is thymidylate. A prominent example for a nucleoside (thymidine) analog is 3'-azido-3'-deoxythymidine (AZT) used in the treatment of AIDS.

The term "kinase activity for a nucleoside or nucleotide analog", within the meaning of the present invention refers to the capability of an enzyme to catalyze the phosphorylation of a nucleoside or nucleotide analog to its corresponding mono-, di- or triphosphate. Naturally, such enzymes, also termed kinases herein, transfer a phosphoryl group from a donor molecule (usually ATP) to an acceptor molecule (in the present invention a nucleoside, nucleotide monophosphate, or a nucleotide diphosphate). In the case of thymidylate kinase, the physiological substrates are ATP and dTMP resulting in ADP and dTDP. AZT is very similar to dTMP but has an azido moiety at the 3' position instead of a hydroxyl group, and is thus also activated by the same enzymes that activate thymidine to thymidine triphosphate (dTTP, a substrate for DNA polymerases).

The term "P-loop", as used herein means a motif that has been identified in many ATP- and GTP-binding proteins (Saraste et al., 1990). The binding of the nucleoside or nucleotide to the P-loop is through main-chain nitrogen atoms to the phosphates of the nucleotide, and through a strictly conserved lysine. A comparison of the amino acid sequences of, for example, thymidylate kinases reveal the consensus sequence recited in (a) above. However, other kinases display this motif as well and could thus be used in accordance of the present invention.

The term "LID region" refers to the mobile part of the kinase enzyme that in case of so called class II (type II) TmpKs (bacterial TmpKs such as from *E. coli*), carries one or more catalytically important arginine or lysine residues and undergoes substantial structural rearrangement upon substrate binding. Accordingly, the LID region of class I (type I) TmpKs (eukaryotic enzymes such as human or yeast

TmpK) constitutes the same flexible region without the catalytically important arginine or lysine residues. As will be appreciated by the person skilled in the art from the disclosure herein the P-loop and/or the LID regions of class II (type II) and class I (type I) enzymes can be transferred into each other via, e.g. amino acid substitution(s), additions(s) and/or deletion(s). Thus, according to the method of the present invention, any protein having nucleoside or nucleotide kinase activity can be used and modified so as to obtain a polypeptide which displays said kinase activity for a nucleoside or nucleotide analog. Moreover, if said protein already displays such activity at a basal level, it is possible to improve its kinase activity for nucleoside and nucleotide analogs compared to the wildtype protein. Furthermore, it is possible using the teaching of the present invention to confer to a polypeptide kinase activity for a nucleoside and nucleotide analog via, e.g., protein design using, for example, computer based redesign of proteins. The resultant proteins and polypeptides obtainable by the method of the present invention can then be tested for their kinase activity by using methods known in the art as described, e.g., in the appended examples or in Guettari (1997).

The present invention is based on the following observations. AZT is a prodrug widely used in the treatment of HIV infection. While the concentration of the active form of AZT, AZT triphosphate (AZT-TTP), reaches only the low micromolar range in human cells exposed to AZT, AZT monophosphate (AZT-MP) accumulates to millimolar concentration. Thus, not only are suboptimal concentrations of the active form of the drug produced, but as a further consequence, a highly toxic intermediate (AZT-MP) is produced at high concentration. The reason behind this accumulation of AZT-MP is the poor phosphorylation of AZT-MP to AZT-DP by the enzyme thymidylate kinase (Furman, Proc. Natl. Acad. Sci. USA 83 (1986), 8333-8337).

In accordance with the present invention, surprisingly it has been found that the reasons behind this low activity of the enzyme is to be found in the interplay of two motifs, the P-loop and the LID-region. In experiments performed in accordance with the present invention, the crystal structure of yeast thymidylate kinase (TmpK)

complexed with the bisubstrate inhibitor P1-(5'-adenosyl) P5-(5'-thymidyl) pentaphosphate (TP₅A) was determined at 2.0 Å resolution. In this complex, TmpK adopts a closed conformation with a region (LID) of the protein closing upon the substrate and forming a helix. The interactions of TmpK and TP₅A revealed that arginine 15, which is located in the phosphate binding loop (P-loop) sequence, plays a catalytic role by interacting with an oxygen atom of the transferred phosphoryl group. Unlike other nucleoside monophosphate kinases where basic residues from the LID region participate in stabilizing the transition state, class I (type I) TmpK lack such residues in the LID region. The present inventors attribute this function to Arg15 of the P-loop. TmpK plays an important role in the phosphorylation of the AIDS prodrug AZT. The structures of TmpK with dTMP and with AZT-MP (Lavie et al, Nature Structural Biology 4 (1997), 601-604) implicate the movement of the Arg15 in response to AZT-MP binding as an important factor for the 200 fold reduced catalytic rate with AZT-MP. TmpK from *E. coli* lacks this arginine in its P-loop while having basic residues in the LID region. This suggested that if such a P-loop movement were to occur in the *E. coli* TmpK upon AZT-MP binding, it should not have such a detrimental effect on catalysis. This hypothesis was tested and as expected by the inventors *E. coli* TmpK phosphorylates AZT-MP only 2.5 times slower than dTMP.

Based on the understanding gained from the present work it became possible to mutate the wild type enzyme in order to attain higher activity for AZT-MP. With the advent of gene therapy, it is now conceivable to transfer the gene of such a modified kinase to the HIV-susceptible cells in an infected individual. The subsequent administration of AZT should result in a higher therapeutic index compared with an individual possessing only the wild type kinase.

The next few paragraphs will further illustrate the invention by way of example for TmpKs and describe how kinases catalyze the phosphoryl transfer reaction, and list the possible changes the person skilled in the art can consider in accordance with the present invention, to achieve (higher) kinase activity for nucleoside or nucleotide analogs.

Kinases transfer a phosphoryl group from a donor molecule (usually ATP) to an acceptor molecule (in the present invention a nucleoside, nucleotide monophosphate, or a nucleotide diphosphate). In the case of thymidylate kinase, the physiological substrates are ATP and dTMP resulting in ADP and dTDP. AZT is very similar to dTMP but has an azido moiety at the 3' position instead of an hydroxyl group, and is thus also activated by the same enzymes that activate thymidine to thymidine triphosphate (dTTP, a substrate for DNA polymerases). Enzymes catalyze chemical reactions by preferentially stabilizing the transition state of a reaction over the corresponding ground state. Since it is highly probable that in phosphoryl transfer reactions an additional negative charge is formed at the transition state in comparison to the total charge of the ground state, stabilizing such a charge lowers the energy barrier for the reaction. Enzymes can stabilize negative charges either by using positively charged amino acid residues (Arg, Lys, or His) or by binding metals, or both. Kinases such as thymidylate kinases utilize both of these possibilities, but since the interaction with the metal (a bound magnesium ion) is probably similar in the ground and transition states, the energy barrier between ground and transition state is not appreciably decreased (this does not mean that the metal is not important for catalysis; the absence of metal abolishes all activity). It is the positive side chain of arginine residue(s) that change their degree of interaction between the ground and transition states and thus achieve preferential stabilization of the transition state over the ground state (i.e. a lowering of the energy barrier for the phosphoryl transfer).

The structure of thymidylate kinase complexed with the bisubstrate analog TP₅A that has been solved in accordance with the present invention allows to observe which basic residues interact with the nucleotides. To date a unique feature of class I (type I) TmpKs is that an arginine from the P-loop interacts directly with the transferred phosphoryl group. The present-inventors postulate that this interaction is responsible for the low activity of TmpK with AZT-MP since the azido group of AZT pushes the P-loop from its position in comparison to its position when dTMP is bound.

An important observation made in accordance with the present invention is that the *E. coli* TmpK (a class II (type II) enzyme) phosphorylates AZT-MP very rapidly. The *E. coli* TmpK has no arginine in its P-loop (a glycine instead) but rather a number of basic residues in a structural motif that is called LID region. This is an example of the scaffold nature of enzymes: Class I (type I) TmpKs have a basic residue in the P-loop motif but none in the LID, whereas class II (type II) TmpKs have the reverse; i.e. none in the P-loop and a few in the LID. In other words, class I (type I) kinases use the P-loop arginine to stabilize the transition state whereas class II (type II) use arginine(s) from the LID for the same purpose.

The above result has immediate implications. Instead of modifying the human or other class I (type I) kinase to better phosphorylate AZT-MP, the *E. coli* TmpK could be used for this purpose (and possibly any other class II (type II) TmpK). The kinetic results described in the appended examples demonstrate that the *E. coli* TmpK is at least 300-fold faster in phosphorylating AZT-MP than the human TmpK. However, because of immunological and other reasons it might be advantageous to use a modified human TmpK over the *E. coli* TmpK in the proposed gene therapy procedure. The modification of the human (or any other class I (type I)) TmpK would be according to the method invention as described above. Advantageously the method of the invention comprises at least one of the following steps:

- Mutation of the LID region of class I (type I) thymidylate kinases to contain one or more basic residues and in doing so mimic the LID motif of class II (type II) TmpKs. This mutation of the LID could be with or without a concomitant mutation in the P-loop. A concomitant substitution in the P-loop (arginine to a smaller amino acid) might be advantageous in order to avoid a steric clash between the P-loop basic residue and the newly introduced LID basic residue(s). The results obtained in accordance with the present invention demonstrate that such an approach is appropriate but they do not rule out that, for example, a lysine in the P-loop might be tolerated.

- Substitution of the entire LID of a class I (type I) TmpK with that from a class II (type II) TmpK, with or without a concomitant mutation in the P-loop; see, e.g., Example 8.
- Selection of appropriate mutant enzymes by measurements of the catalytic efficiency with a coupled colorimetric assay that links the phosphorylation of, e.g., the nucleoside monophosphate to diphosphate to a change of the optical density. This can be extended to a high throughput screening (HTS) setup with a microtiter plate reader.


The method of the invention can be performed using conventional techniques known in the art, for example, by using amino acid deletion(s), insertion(s), substitution(s), addition(s), and/or recombination(s) and/or any other modification(s) known in the art either alone or in combination. Methods for introducing such modifications in the DNA sequence underlying the amino acid sequence of a protein having kinase activity are well known to the person skilled in the art; see, e.g., Sambrook, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y.

In a preferred embodiment of the method of the invention said protein to be modified with the method described above is derived from a eukaryotic or prokaryotic organism, preferably said organism is human or yeast. As described above, the present inventors have solved the 3-dimensional structure of TmpK complexed with the bisubstrate inhibitor P1-(5'-Adenosyl)P5-(5'-Thymidyl) pentaphosphate (TP₅A). This structure, taken together with those of TmpK complexed with dTMP and with AZT-MP (Lavie et al., 1997b) allowed the inventors to identify the residues important for catalysis, and to propose an explanation for the slow phosphorylation of AZT-MP. In addition, based on that knowledge, it permits the prediction that the rate of AZT-MP phosphorylation by the *E. coli* TmpK would be similar to the phosphorylation rate of the physiological substrate dTMP. In fact, as shown in accordance with the present invention *E. coli* TmpK discriminates poorly against AZT-MP, supporting the inventors finding that Arg15

of yeast and human thymidylate kinases is responsible for discrimination against AZT-MP as a substrate (the human thymidylate kinase has 44% amino acid identity and >63% similarity with the yeast enzyme with all catalytically important residues conserved between the two species; see Figure 4). Furthermore, as demonstrated in Example 8 herein below the yeast TmpK could, by applying the method of the invention, converted into an enzyme displaying high kinase activity for the nucleoside analog AZT. Similar results are to be expected when using the human or any other corresponding eukaryotic enzyme.

In a preferred embodiment of the method of the invention said protein comprises the amino acid sequence of any one of SEQ ID NOS: 1 to 13 or a fragment thereof. The amino acid sequences depicted in SEQ ID NOS: 1 to 13 belong to various TmpKs of eukaryotic and prokaryotic origin and comprise, for example, the amino acid sequence of *E. coli* TmpK. With the method of the present invention and the teaching provided herein, it is, e.g., possible to improve the kinase activity of the human or yeast enzyme for nucleoside or nucleotide analogs, in particular for AZT. Furthermore, the teaching of the present invention now enables rationale shuffling or domains from, e.g., *E. coli* TmpK to the corresponding enzyme from human, mouse or yeast. On the other hand it is also possible to modify the amino acid sequence of the *E. coli* enzyme to more closely resemble that of the corresponding eukaryotic, preferably human enzyme while the P-loop and the LID region remain substantially unaffected and therefore the resultant polypeptide retains its kinase activity for nucleoside and nucleotide analogs. For the rational design of polypeptides produced according to the method of the invention computer programs may be used such as BRASMOL that are obtainable from the Internet. Furthermore, folding simulations and computer redesign of structural motifs can be performed using other appropriate computer programs (Olszewski, Proteins 25 (1996), 286-299; Hoffman, Comput. Appl. Biosci. 11 (1995), 675-679). Computers can be used for the conformational and energetic analysis of detailed protein models (Monge, J. Mol. Biol. 247 (1995), 995-1012; Renouf, Adv. Exp. Med. Biol. 376 (1995), 37-45).

In another preferred embodiment the method of the present invention said amino acid which is substituted or added in (a) is glycine or lysine. In case that the amino acid is changed in the P-loop of class I (type I) enzymes the amino acid that is changed is preferably a small amino acid such as glycine. However, alanine, less preferably valine, threonine, serine or glutamic acid may work as well although probably less effective. As described above, the LID region of class I (type I) thymidylate kinases may be substituted in accordance with the method of the invention to contain at least one basic residue. Thus, in a preferred embodiment of the method of the invention said amino acid which is substituted or added in (b) is a basic amino acid, preferably arginine. Usually, said LID has the consensus sequence R/KXXXXXERYEXXXQ. This consensus sequence or substantially identical sequences can be determined by comparison of the amino acid sequences of known nucleoside and nucleotide kinases, for example those shown in SEQ ID NOS: 1 to 13. Preferably, said polypeptide obtainable by the method of the invention exhibits kinase activity for a nucleoside or nucleotide analog which is higher than that of the corresponding wild type enzyme, preferably higher than that of an eukaryotic, most preferably higher than that of the human enzyme. Preferably, said kinase activity for a nucleoside or nucleotide analog is 5-fold, more preferably 10-fold, still more preferably 30-fold and most preferably 300-fold improved compared to the corresponding wild-type enzyme. Furthermore, it is preferred within the method of the invention that said nucleoside analog is AZT, d4T or has the following structure:



wherein B is any nucleobase or analog thereof, and X is O, CH₂, NH or S.

As discussed above, the amino acid substitution(s) or other modifications performed in the amino acid sequence of a given kinase may result in the P-loop and/or in the LID region of a bacterial nucleoside or nucleotide kinase, preferably those of the TmpK of *E.coli*; see also Example 8.

As mentioned before, for the long term aim of gene therapeutic potentiation of AZT effectiveness, there could be significant advantages from use of a modified human enzyme. Therefore, the mutational studies were extended to the human enzyme; see Example 9. Contrary to all expectations arising from the studies on the yeast enzyme, replacement of Arg-16 (equivalent to Arg-15 in the yeast enzyme) did not lead to a loss in catalytic activity, but to a slight gain. It is possible that this is related to the fact that the k_{cat} for the human enzyme is much slower than for the yeast enzyme (0.67 cf. 35 s⁻¹), which is already a very slow kinase when compared with other nucleoside monophosphate kinases. Without intending to be bound by theory it is believed that since the catalytic machinery in both enzymes appears to be identical, as shown by sequence comparison and 3-D structural determination, it is possible that the chemical step (i.e. phosphate transfer) is not rate limiting, but rather product release, so that slowing down a relatively rapid chemical step might not have any influence on the overall rate.

In keeping with expectations arising from the yeast TMPK data was the fact that introduction of the *E. coli* lid region without removing Arg-16 led to a drop in catalytic activity. However, and most dramatically, a combination of introduction of the *E. coli* lid with replacement of Arg-16 by glycine not only restored more than full wild type catalytic activity, but resulted in a protein which is more efficient with AZT-MP than with TMP. The increase in activity with AZT-MP is approximately 300 fold. This mutant thus has properties which are highly attractive for improving the potency of AZT, since the efficiency of AZTMP phosphorylation is improved dramatically with only a minor increase in TMP phosphorylation activity. The latter is an important aspect, since AZTTP must compete with TTP for HIV-reverse transcriptase catalysed addition to the end of a growing (HIV) DNA chain.

The mutants of TMPK described so far showing altered specificity for AZTMP and TMP were produced according to rational considerations based on comparative structure-function studies of TMPK from three different sources. In this respect, a further striking result is that obtained by replacing Phe-105 in the human enzyme by tyrosine. The rational for this was that in the yeast enzyme, the corresponding residue (at position 102) is a tyrosine, and its hydroxyl group interacts with the carboxylate side chain of Asp-14, which (like Asp-15 in the human enzyme) appears to be an essential residue for the catalytic mechanism. The Phe-105 Tyr mutant of the human enzyme shows reduced activity with TMP, but, unexpectedly, significantly increased activity with AZTMP, so that the latter is now a better substrate than the former; see Example 9. Thus, and again without intending to be bound by theory it is believed that substitution of amino acid(s) that interact with the LID or P-loop, and in particular, if present, with the carboxylate side chain of Asp leads to a conformational change of the catalytic center of the kinase enzyme and fulfils the requirement of greatly increased AZTMP phosphorylating activity without increasing (actually decreasing) TMP phosphorylation. An amino acid position in a nucleotide or nucleoside kinase corresponding to amino acid residue 105 in the human thymidylate kinase can be determined by comparison and alignment of the amino acid sequences of known nucleoside and nucleotide kinases, for example, those shown in SEQ ID NOS: 1 to 13 using, e.g., the program GCG. The specificity ratio (AZTMP:TMP) is actually identical with that of the mutant containing Gly-16 and the E. coli lid, although the overall activity is a factor of ca. 10 lower.

The results obtained in accordance with the present invention show how rational considerations have led to generation of TMPK mutants capable of phosphorylating the nucleoside pro-drug AZT with high efficiency. This finding gives rise to several applications in the medical and diagnostic field which will be explained in more detail below.

In another embodiment the present invention relates to a polynucleotide encoding the polypeptide obtainable by the method of the invention. Said polynucleotide

may be, e.g., DNA, cDNA, genomic DNA, RNA or synthetically produced DNA or RNA or a recombinantly produced chimeric nucleic acid molecule comprising any of those polynucleotides either alone or in combination. Preferably said polynucleotide is part of a vector. Such vectors may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions.

In a further preferred embodiment the polynucleotide of the invention is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic cells.

Expression of said polynucleotide comprises transcription of the polynucleotide into a translatable mRNA. Regulatory elements ensuring expression in eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art. They usually comprise regulatory sequences ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the *lac*, *trp* or *tac* promoter in *E. coli*, and examples for regulatory elements permitting expression in eukaryotic host cells are the *AOX1* or *GAL1* promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (In-vitrogene); pSPORT1 (GIBCO BRL).

The polynucleotide of the invention can be used alone or as part of a vector to express the polypeptide having kinase activity for nucleoside or nucleotide analogs in cells, for, e.g., gene therapy or diagnostics of diseases related to viral infections

and cancer. The polynucleotide or vector containing the DNA sequence encoding a kinase for nucleoside or nucleotide analogs is introduced into the cells which in turn produce the protein of interest. Gene therapy, which is based on introducing therapeutic genes into cells by *ex-vivo* or *in-vivo* techniques is one of the most important applications of gene transfer. Suitable vectors and methods for *in-vitro* or *in-vivo* gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, Nature Medicine 2 (1996), 534-539; Schaper, Circ. Res. 79 (1996), 911-919; Anderson, Science 256 (1992), 808-813; Isner, Lancet 348 (1996), 370-374; Muhlhauser, Circ. Res. 77 (1995), 1077-1086; Wang, Nature Medicine 2 (1996), 714-716; WO94/29469; WO 97/00957 or Schaper, Current Opinion in Biotechnology 7 (1996), 635-640, and references cited therein. The polynucleotides and vectors of the invention may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Preferably, said cell is a germ line cell, embryonic cell, or egg cell or derived therefrom, most preferably said cell is a stem cell.

Furthermore, the present invention relates to vectors, particularly plasmids, cosmids, viruses and bacteriophages used conventionally in genetic engineering that comprise a polynucleotide of the invention. Preferably, said vector is an expression vector and/or a gene transfer or targeting vector. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the polynucleotides or vector of the invention into targeted cell population. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1989). Alternatively, the polynucleotides and vectors of the invention can be reconstituted into liposomes for delivery to target cells.

The present invention furthermore relates to host cells transformed with a polynucleotide or vector of the invention. Said host cell may be a prokaryotic or eukaryotic cell. The polynucleotide or vector of the invention which is present in the host cell may either be integrated into the genome of the host cell or it may be maintained extrachromosomally. In this respect, it is also to be understood that the recombinant DNA molecule of the invention can be used for "gene targeting" and/or "gene replacement", for restoring a mutant gene or for creating a mutant gene via homologous recombination; see for example Mouellic, Proc. Natl. Acad. Sci. USA, 87 (1990), 4712-4716; Joyner, Gene Targeting, A Practical Approach, Oxford University Press.

The host cell can be any prokaryotic or eukaryotic cell, such as a bacterial, insect, fungal, plant, animal or human cell. Preferred fungal cells are, for example, those of the genus *Saccharomyces*, in particular those of the species *S. cerevisiae*. The term "prokaryotic" is meant to include all bacteria which can be transformed or transfected with a polynucleotide for the expression of a polypeptide of the invention. Prokaryotic hosts may include gram negative as well as gram positive bacteria such as, for example, *E. coli*, *S. typhimurium*, *Serratia marcescens* and *Bacillus subtilis*. A polynucleotide coding for a polypeptide of the invention can be used to transform or transfect the host using any of the techniques commonly known to those of ordinary skill in the art. Especially preferred is the use of a plasmid or a virus containing the coding sequence of the polypeptide having kinase activity for a nucleoside or nucleotide analog for purposes of prokaryotic transformation or transfection, respectively. Methods for preparing fused, operably linked genes and expressing them in bacteria are well-known in the art (Maniatis, et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1989). The genetic constructs and methods described therein can be utilized for expression of the polypeptide of the invention in prokaryotic hosts. In general, expression vectors containing promoter sequences which facilitate the efficient transcription of the inserted polynucleotide are used in connection with the host. The expression vector typically contains an origin of replication, a promoter, and a terminator, as well as specific genes which are

capable of providing phenotypic selection of the transformed cells. The transformed prokaryotic hosts can be grown in fermentors and cultured according to techniques known in the art to achieve optimal cell growth. The polypeptides of the invention can then be isolated from the grown medium, cellular lysates, or cellular membrane fractions. The isolation and purification of the microbially expressed polypeptides of the invention may be by any conventional means such as, for example, preparative chromatographic separations and immunological separations such as those involving the use of monoclonal or polyclonal antibodies.

Thus, in a further embodiment the invention relates to a method for the production of a polypeptide having kinase activity for a nucleoside or nucleotide analog comprising culturing a host cell as defined above under conditions allowing the expression of the polypeptide and recovering the produced polypeptide from the culture.

In another embodiment the present invention relates to a method for producing cells capable of expressing a polypeptide having nucleoside or nucleotide kinase activity for nucleoside or nucleotide analogs comprising genetically engineering cells with the polynucleotide or with the vector of the invention. The cells obtainable by the method of the invention can be used, for example, to test the capacity of the polypeptides of the invention to improve the metabolism of nucleoside or nucleotide analogs and, if mammalian cells are used, to test their impact on anti-viral activity of nucleoside and nucleotide analogs. Furthermore, the cells can be used to study known and unknown nucleoside and nucleotide analogs for their ability to be converted to the corresponding mono-, di- or triphosphates. The cells obtainable by the above-described method may also be used for the screening methods referred to herein below.

Furthermore, the invention relates to a polypeptide having kinase activity for a nucleoside or nucleotide analog encoded by a polynucleotide according to the

invention or obtainable by the above-described methods or from cells produced by the method described above.

In this context it is also understood that the polypeptides according to the invention may be further modified by conventional methods known in the art. By providing the polypeptides according to the present invention it is also possible to determine the portions relevant for their biological activity, namely their kinase activity. This may allow the construction of chimeric proteins comprising an amino acid sequence derived from a polypeptide of the invention which is crucial for kinase activity and other functional amino acid sequences e.g. nuclear localization signals, transactivating domains, DNA-binding domains, hormone-binding domains, protein tags (GST, GFP, h-myc peptide, Flag, HA peptide) which may be derived from the same or from heterologous proteins.

The present invention furthermore relates to antibodies specifically recognizing a polypeptide according to the invention which has kinase activity for a nucleoside or nucleotide analog. Advantageously, the antibody specifically recognizes a polypeptide according to the invention which has kinase activity for a nucleoside or nucleotide analog but does not recognize a polypeptide which is a wild type starting protein of such a polypeptide and which has no or less kinase activity for nucleoside or nucleotide analogs than the polypeptides of the invention.

Antibodies against the polypeptide of the invention can be prepared by well known methods using a purified polypeptide according to the invention or a synthetic fragment derived therefrom as an antigen. Monoclonal antibodies can be prepared, for example, by the techniques as originally described in Köhler and Milstein, Nature 256 (1975), 495, and Galfré, Meth. Enzymol. 73 (1981), 3, which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals. The antibodies can be monoclonal antibodies, polyclonal antibodies or synthetic antibodies as well as fragments of antibodies, such as Fab, Fv or scFv fragments etc. Furthermore, antibodies or fragments thereof to the aforementioned polypeptides can be obtained by using methods which are

described, e.g., in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. These antibodies can be used, for example, for the immunoprecipitation and immunolocalization of the polypeptides of the invention as well as for the monitoring of the presence of such polypeptides, for example, in recombinant organisms, and for the identification of compounds interacting with the proteins according to the invention. For example, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies which bind to an epitope of the polypeptide of the invention (Schier, Human Antibodies Hybridomas 7 (1996), 97-105; Malmberg, J. Immunol. Methods 183 (1995), 7-13).

Moreover, the present invention relates to a composition, preferably a pharmaceutical composition comprising

- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide; or
- (b) the above-described polypeptide, polynucleotide or vector of the invention;
- (c) optionally a nucleoside or nucleotide analog; and
- (d) optionally a pharmaceutically acceptable carrier.

As described above and as shown in the appended examples prokaryotic kinases such as *E. coli* TmpK belong to class II (type II) enzymes which display superior phosphorylation properties for nucleoside and nucleotide analogs, in particular for AZT. Thus, it is not necessary to further modify such prokaryotic proteins according to the method of the invention but it is possible to employ them in a pharmaceutical composition of the invention unmodified or substantially unmodified.

Examples of suitable pharmaceutical carriers are well known in the art and include phosphate buffered saline solutions, water, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions etc. Compositions comprising such carriers can be formulated by well known conventional methods.

These pharmaceutical compositions can be administered to the subject at a suitable dose. Administration of the suitable compositions may be effected by different ways, e.g., by intravenous, intraperitoneal, subcutaneous, intramuscular, topical or intradermal administration. The dosage regimen will be determined by the attending physician and other clinical factors. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Generally, the regimen as a regular administration of the pharmaceutical composition should be in the range of 1 μ g to 10 mg units per day. If the regimen is a continuous infusion, it should also be in the range of 1 μ g to 10 mg units per kilogram of body weight per minute, respectively. Progress can be monitored by periodic assessment. Dosages will vary but a preferred dosage for intravenous administration of DNA is from approximately 10^6 to 10^{12} copies of the DNA molecule. The compositions of the invention may be administered locally or systemically. Administration will generally be parenterally, e.g., intravenously; DNA may also be administered directly to the target site, e.g., by biolistic delivery to an internal or external target site or by catheter to a site in an artery.

It is envisaged by the present invention that the various polynucleotides and vectors of the invention are administered either alone or in any combination using standard vectors and/or gene delivery systems, and optionally together with an appropriate compound, for example a nucleoside or nucleotide analog, and/or together with a pharmaceutically acceptable carrier or excipient. Subsequent to administration, said polynucleotides or vectors may be stably integrated into the genome of the subject. On the other hand, viral vectors may be used which are specific for certain cells or tissues, preferably for CD4 cells and persist in said cells. Suitable pharmaceutical carriers and excipients are well known in the art. The pharmaceutical compositions prepared according to the invention can be used for the prevention or treatment or delaying of different kinds of diseases, which are related to viral infection or cancer.

Furthermore, it is possible to use a pharmaceutical composition of the invention which comprises polynucleotide or vector of the invention in gene therapy. Suitable gene delivery systems may include liposomes, receptor-mediated delivery systems, naked DNA, and viral vectors such as herpes viruses, retroviruses, adenoviruses, and adeno-associated viruses, among others. Delivery of nucleic acids to a specific site in the body for gene therapy may also be accomplished using a biolistic delivery system, such as that described by Williams (Proc. Natl. Acad. Sci. USA 88 (1991), 2726-2729).

Standard methods for transfecting cells with recombinant DNA are well known to those skilled in the art of molecular biology, see, e.g., WO 94/29469. Gene therapy may be carried out by directly administering the recombinant DNA molecule or vector of the invention to a patient or by transfecting cells such as CD4 with the polynucleotide or vector of the invention ex vivo and infusing the transfected cells into the patient. Furthermore, research pertaining to gene transfer into cells of the germ line is one of the fastest growing fields in reproductive biology. Gene therapy, which is based on introducing therapeutic genes into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable vectors and methods for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., WO 94/29469, WO 97/00957 or Schaper (Current Opinion in Biotechnology 7 (1996), 635-640) and references cited above. The polynucleotides and vectors comprised in the pharmaceutical composition of the invention may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) containing said recombinant DNA molecule into the cell. Preferably, said cell is a germ line cell, embryonic cell, stem cell or egg cell or derived therefrom. The pharmaceutical compositions according to the invention can be used for the treatment of diseases hitherto unknown as being related to viral infection or cancer. An embryonic cell can be for example an embryonic stem cell as described in, e.g., Nagy, Proc. Natl. Acad. Sci. USA 90 (1993) 8424-8428.

It is to be understood that the introduced polynucleotides and vectors of the invention express the polypeptide or protein having kinase activity for nucleoside or nucleotide analogs after introduction into said cell and preferably remain in this status during the lifetime of said cell. For example, cell lines which stably express the polynucleotide under the control of appropriate regulatory sequences may be engineered according to methods well known to those skilled in the art. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with the polynucleotide or vector of the invention and a selectable marker, either on the same or separate vectors. Following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows for the selection of cells having stably integrated the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. Such engineered cell lines are particularly useful in screening methods described below.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, Cell 11(1977), 223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska, Proc. Natl. Acad. Sci. USA 48 (1962), 2026), and adenine phosphoribosyltransferase (Lowy, Cell 22 (1980), 817) in tk⁻, hgprt⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler, Proc. Natl. Acad. Sci. USA 77 (1980), 3567; O'Hare, Proc. Natl. Acad. Sci. USA 78 (1981), 1527), gpt, which confers resistance to mycophenolic acid (Mulligan, Proc. Natl. Acad. Sci. USA 78 (1981), 2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, J. Mol. Biol. 150 (1981), 1); hygromycin, which confers resistance to hygromycin (Santerre, Gene 30 (1984), 147); or puromycin (pat, puromycin N-acetyl transferase). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman, Proc. Natl. Acad. Sci. USA 85 (1988), 8047); and ODC (ornithine decarboxylase)

which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue, 1987, In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed.). Cells to be used for *ex vivo* gene therapy are well known to those skilled in the art. For example, mature cells of the immune system present in blood or preferably the corresponding stem cells.

In a preferred embodiment of the invention said protein in the pharmaceutical composition is a bacterial nucleoside or nucleotide kinase, preferably a bacterial TmpK. As described in the appended examples the TmpK of *E. coli* or its structure with respect to the P-loop and the LID region appear to be particularly suited for the activation of nucleoside and nucleotide analogs, e.g., AZT. Thus, in a preferred embodiment of the pharmaceutical composition of the invention said protein to be employed in accordance with the present invention has at least the P-loop and/or the LID region of *E. coli* TmpK, for example, said TmpK comprises the amino acid sequence of *E. coli* TmpK, e.g., that shown in SEQ ID NO: 4 or a biologically active fragment thereof.

The present invention also relates to compositions comprising at least one of the aforementioned polynucleotides, vectors, polypeptides, proteins or antibodies, and in the case of kits or diagnostic compositions, optionally suitable means for detection. Said compositions may further contain compounds such as nucleoside or nucleotide analogs, further plasmids, antibiotics and the like for screening transgenic cells useful for the genetic engineering of non-human animals, preferably mammals and most preferably mouse. The diagnostic compositions of the invention may be used for methods of detecting and isolating nucleoside or nucleotide analogs which are functionally equivalent to, e.g., AZT or d4T.

The present invention also relates to a method for the production of a transgenic non-human animal, preferably transgenic mouse, comprising introduction of a polynucleotide or vector of the invention into a germ cell, an embryonic cell, stem

cell or an egg or a cell derived therefrom. The non-human animal to be used in the method of the invention may be a non-transgenic healthy animal, or may have a viral disease or cancer, preferably a disease caused by infection of a retrovirus such as HIV, HTLV or related viruses. Production of transgenic embryos and screening of those can be performed, e.g., as described by A. L. Joyner Ed., Gene Targeting, A Practical Approach (1993), Oxford University Press. The DNA of the embryonal membranes of embryos can be analyzed using Southern blots with an appropriate probe.

The invention also relates to transgenic non-human animals such as transgenic non-human mouse, rats, hamsters, dogs, monkeys, rabbits or pigs comprising a polynucleotide or vector of the invention or obtained by the method described above, preferably wherein said polynucleotide or vector is stably integrated into the genome of said non-human animal, preferably such that the presence of said polynucleotide or vector leads to the expression of the polypeptide of the invention.

With the polypeptides, polynucleotides and vectors of the invention, it is now possible to study *in vivo* and *in vitro* the efficiency of nucleoside and nucleotide analogs. Furthermore, since the polypeptides of the invention provide for optimal or at least improved activation of said nucleoside or nucleotide analog, it is now possible to determine further analogs which may be effective for the treatment of viral diseases or cancer, for example specific tumors or AIDS.

The present invention further relates to a method for identifying an inhibitor of a nucleoside or nucleotide kinase comprising the steps of:

- (a) contacting the polypeptide of the invention or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a compound to be screened under conditions that permit binding of said compound to the nucleoside or nucleotide kinase, and

- (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the absence or decrease of the signal is indicative for an inhibitor of a nucleoside or nucleotide kinase.

Furthermore, the invention relates to a method for identifying a nucleoside or nucleotide based prodrug comprising the steps of

- (a) contacting the polypeptide of the invention or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a nucleoside or nucleotide analog compound to be screened under conditions that permit kinase activity of said polypeptide, and
- (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the presence of a signal is indicative for a putative prodrug.

The term "compound" in a method of the invention includes a single substance or a plurality of substances which may or may not be identical.

Said compound(s) may be comprised in, for example, samples, e.g., cell extracts from, e.g., plants, animals or microorganisms. Furthermore, said compounds may be known in the art but hitherto not known to be capable of inhibiting a nucleoside or nucleotide kinase or not known to be useful as a prodrug, respectively. The plurality of compounds may be, e.g., added to the culture medium or injected into a cell or non-human animal of the invention.

If a sample containing (a) compound(s) is identified in the method of the invention, then it is either possible to isolate the compound from the original sample identified as containing the compound, in question or one can further subdivide the original sample, for example, if it consists of a plurality of different compounds, so as to reduce the number of different substances per sample and repeat the method with the subdivisions of the original sample. It can then be determined whether said sample or compound displays the desired properties, for example, by the methods described herein, the appended examples or in the literature, e.g., Guettari (1997). Depending on the complexity of the samples, the steps described above can be

performed several times, preferably until the sample identified according to the method of the invention only comprises a limited number of or only one substance(s). Preferably said sample comprises substances of similar chemical and/or physical properties, and most preferably said substances are identical. The methods of the present invention can be easily performed and designed by the person skilled in the art, for example in accordance with other cell based assays described in the prior art or by using and modifying the methods as described in the appended examples. Furthermore, the person skilled in the art will readily recognize which further compounds and/or enzymes may be used in order to perform the methods of the invention, for example, enzymes, if necessary, that convert a certain compound into the precursor which in turn represents a substrate for the kinase of the invention. Such adaptation of the method of the invention is well within the skill of the person skilled in the art and can be performed without undue experimentation.

Compounds which can be used in accordance with the present invention include peptides, proteins, nucleic acids, antibodies, small organic compounds, ligands, peptidomimetics, PNAs and the like. Said compounds can also be functional derivatives or analogues of known nucleoside or nucleotide analogs. Methods for the preparation of chemical derivatives and analogues are well known to those skilled in the art and are described in, for example, Beilstein, Handbook of Organic Chemistry, Springer edition New York Inc., 175 Fifth Avenue, New York, N.Y. 10010 U.S.A. and Organic Synthesis, Wiley, New York, USA. Furthermore, said derivatives and analogues can be tested for their effects according to methods known in the art or as described, for example, in the appended examples. Furthermore, peptide mimetics and/or computer aided design of appropriate nucleoside and nucleotide derivatives and analogues can be used, for example, according to the methods described below. Nucleoside and nucleotide analogs comprise molecules having as the basis structure a ribo-, deoxyribo- or dideoxyribonucleoside.

Appropriate computer programs can be used for the identification of interactive sites of a putative inhibitor and the polypeptide of the invention by computer assistant searches for complementary structural motifs (Fassina, Immunomethods 5 (1994), 114-120). Further appropriate computer systems for the computer aided design of protein and peptides are described in the prior art, for example, in Berry, Biochem. Soc. Trans. 22 (1994), 1033-1036; Wodak, Ann. N. Y. Acad. Sci. 501 (1987), 1-13; Pabo, Biochemistry 25 (1986), 5987-5991. The results obtained from the above-described computer analysis can be used in combination with the method of the invention for, e.g., optimizing known nucleoside or nucleotide analogs. Appropriate peptide mimetics and other inhibitors can also be identified by the synthesis of peptide mimetic combinatorial libraries through successive chemical modification and testing the resulting compounds, e.g., according to the methods described herein and in the appended examples. Methods for the generation and use of peptide mimetic combinatorial libraries are described in the prior art, for example in Ostresh, Methods in Enzymology 267 (1996), 220-234 and Dorner, Bioorg. Med. Chem. 4 (1996), 709-715. Furthermore, the three-dimensional and/or crystallographic structure of inhibitors of the polypeptide of the invention can be used for the design of peptide mimetic inhibitors or nucleoside and nucleotide analogs, e.g., in combination with the polypeptide of the invention (Rose, Biochemistry 35 (1996), 12933-12944; Rutenber, Bioorg. Med. Chem. 4 (1996), 1545-1558).

The compounds identified according to the method of the invention, in particular nucleoside and nucleotide analogs are expected to be very beneficial since the prodrugs that have been used so far are only of limited use due to their inefficient metabolism in the subject.

In summary, the present invention provides methods for identifying compounds which inhibit kinase activity as well as compounds that can be used as nucleoside or nucleotide based prodrugs even in the absence of a polypeptide or protein having improved kinase activity for nucleoside and nucleotide analogs.

Compounds found to downregulate the activity of kinases may be used in the treatment of cancer and related diseases. In addition, it may also be possible to specifically inhibit viral kinases, thereby preventing viral infection or viral spread.

The compounds identified or obtained according to the method of the present invention are thus expected to be very useful in diagnostic and in particular for therapeutic applications. Hence, in a further embodiment the invention relates to a method for the production of a pharmaceutical composition comprising the steps of

- (a) contacting the polypeptide of the invention or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a compound to be screened under conditions that permit binding of said compound to the nucleoside or nucleotide kinase, and
- (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the absence or decrease of the signal is indicative for an inhibitor of a nucleoside or nucleotide kinase, or
- (a') contacting the polypeptide of the invention or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a nucleoside or nucleotide analog compound to be screened under conditions that permit kinase activity of said polypeptide, and
- (b') detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the presence of a signal is indicative for a putative prodrug; and
- (c) formulating the inhibitor identified in step (b) or the nucleoside or nucleotide analog identified in step (b') in a pharmaceutically acceptable form.

The therapeutically useful compounds identified according to the method of the invention may be administered to a patient by any appropriate method for the particular compound, e.g., orally, intravenously, parenterally, transdermally, transmucosally, or by surgery or implantation (e.g., with the compound being in the form of a solid or semi-solid biologically compatible and resorbable matrix) at or

near the site where the effect of the compound is desired. Therapeutic doses are determined to be appropriate by one skilled in the art, see *supra*.

Furthermore, the present invention relate to the use of

- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide,
- (b) the polypeptide, the polynucleotide or the vector of the invention; and/or
- (c) the nucleoside or nucleotide analog identified in the method of the invention for the preparation of a pharmaceutical composition for the activation of nucleoside or nucleotide analogs or nucleoside or nucleotide based prodrugs and/or for the treatment of viral infections and/or diseases or cancer. Preferably said activation results in a cytotoxic nucleoside or nucleotide. In a preferred embodiment of the use of the invention said viral infection is HIV infection.

In a further embodiment the present invention relates to the use of the inhibitor obtainable by the method of the invention for the preparation of a pharmaceutical composition for inhibiting virus replication or for treating cancer. As mentioned above, the inhibitor identified and obtainable by the method of the invention may be used to specifically inhibit the activity of a viral kinase. This would mean that the activation of nucleosides and/or nucleotides necessary for viral replication is suppressed and therefore may result in preventing of the production of viral progeny.

In a preferred embodiment of the invention, the pharmaceutical compositions, e.g., to be used as described above further comprise or are designed to be administered with a nucleoside or nucleotide analog, preferably AZT or d4T.

In a still further embodiment, the present invention relates to a method for the preparative synthesis of a nucleoside phosphate analog comprising:

- (a) using a polynucleotide of the invention or as defined above in a noncellular system or in a cell *ex vivo*, and
- (b) formulating the cells modified in step (a) in a pharmaceutically acceptable form.

In another embodiment the present invention relates to the use of

- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide, or

- (b) the polypeptide, the polynucleotide or the vector of invention

for the preparation of nucleoside phosphates or analogs and derivatives thereof.

Prior to the present inventions it was cost extensive or even impossible to produce analogs of nucleosides or nucleotides useful in diagnostic and therapeutic approaches; examples are phosphorylated forms of AZT and d4T. With the help of the present invention it is now possible to prepare such nucleoside and nucleotide analogs, for example, β or γ -phosphate labeled nucleotides. For instance, the polypeptide of the invention is particularly suited for use in the preparation of nucleoside or nucleotide analogs in which it can be utilized in liquid phase or bound to a solid phase carrier. The polypeptide of the invention can be bound to many different carriers and used to produce nucleoside and nucleotide analogs. Examples of well-known carriers include glass, polystyrene, polyvinyl chloride, polypropylene, polyethylene, polycarbonate, dextran, nylon, amyloses, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the carrier can be either soluble or insoluble for purposes of the invention. Those skilled in the art will know of other suitable carriers for binding the polypeptide or will be able to ascertain such, using routine experimentation. There are many different labels known to those of ordinary skill in the art. Examples of the types of labels which can be used in the present invention include radioisotopes, colloidal metals, fluorescent compounds, chemiluminescent compounds, and bioluminescent compounds. On the other hand, a cell of the invention may be

used in a fermentation process for the production of such nucleoside or nucleotide analogs.

These and other embodiments are disclosed and encompassed by the description and examples of the present invention. Further literature concerning any one of the methods, uses and compounds to be employed in accordance with the present invention may be retrieved from public libraries and databases, using for example electronic devices. For example the public database "Medline" may be utilized which is available on the Internet, for example under <http://www.ncbi.nlm.nih.gov/PubMed/medline.html>. Further databases and addresses, such as <http://www.ncbi.nlm.nih.gov/>, <http://www.infobiogen.fr/>, http://www.fmi.ch/biology/research_tools.html, <http://www.tigr.org/>, are known to the person skilled in the art and can also be obtained using, e.g., <http://www.lycos.com>. An overview of patent information in biotechnology and a survey of relevant sources of patent information useful for retrospective searching and for current awareness is given in Berks, TIBTECH 12 (1994), 352-364.

The pharmaceutical compositions, uses, methods of the invention can be used advantageously for the treatment of all kinds of diseases hitherto unknown as being related to or dependent on viral diseases or cancer. The pharmaceutical compositions, methods and uses of the present invention may be desirably employed in humans, although animal treatment is also encompassed by the methods and uses described herein.

The figures show:

Figure 1 The TP5A-TmpK complex structure. **a.** A ribbon diagram of a monomer (TmpK is a homodimer) with the helices drawn in red and strands in green. The TP₅A bound at the active site is in cyan with the 5 phosphorous atoms in purple. The LID region adopts a helical conformation in the TP₅A bound complex in contrast to a coil with no

secondary structure when only dTMP present. **b.** Stereo view of the TP₅A-TmpK crystal packing. The monomers can be designated active (green) or inactive (red) according to the presence of a symmetry related arginine (depicted as ball-and-stick) in the active site. The TP₅A molecules are shown in cyan. Note that the arginines in green penetrate the active site of the red monomers (unlike the red-colored arginines that are further away) and thus hinder the formation of an active conformation by those monomers.

Figure 2 **a.** Stereo view of a simulated-annealing omit map of TP₅A and arginine 15 from an active monomer contoured at 3 sigma. Arginine 15, which is located in the P-loop (shown as coil running left-to-right) interacts with the middle phosphate of TP₅A. **b.** Stereo view of the active site. The TP₅A nucleotide is drawn in yellow except for the 5 phosphorous atoms that are in purple. **c.** A distance map with the same view as above. P-loop residues are additionally marked with an asterisk.

Figure 3 Steady-state kinetics of dTMP versus AZT-MP with *E. coli* TmpK.

a. The catalytic activity of *E. coli* TmpK (16.4 nM) with dTMP and ATP was measured as described in Methods at dTMP concentrations indicated at the X-axis, and the following ATP concentrations 15 μ M (\circ), 25 μ M (\bullet), 50 μ M (\square), 100 μ M (\blacksquare), 300 μ M (\triangle) and 1000 μ M (\blacktriangle). The maximal catalytic activity attained was $k_{cat} = 15 \text{ s}^{-1}$ with K_M dTMP of 2.7 μ M with ATP and dTMP binding with a factor for synergism of 6.

b. The catalytic activity of *E. coli* TmpK (15.1 nM) with AZT-MP and ATP. The ATP concentrations were as indicated in a. The turnover number (k_{cat}) reached a maximum of 6 s^{-1} at saturating concentrations of both substrates. The binding of dTMP and ATP is

synergistic by a factor of 6. The apparent K_M of dTMP is lowered from 17 μM (extrapolated to no ATP) to 2.7 μM at saturating ATP concentration, and likewise the apparent K_M value of ATP is lowered from approximately 50 μM to 8 μM . This synergism of binding is completely absent with ATP and AZT-MP, the K_M value of ATP being around 50 μM independent of the AZT-MP concentration and that of AZT-MP (30 μM) being very close to dTMP (17 μM) in the absence of ATP. Therefore, it is predicted that the 3'-OH group of dTMP may be directly involved in the synergism of ATP and dTMP binding.

Figure 4 Sequence alignment (using GCG) of 3 eukaryotic (KTHY_SCHPO is TmpK from *Schizosaccharomyces pombe*) and the *E. coli* thymidylate kinase amino-acid sequences. Shaded black are residues conserved in all sequences, in gray are similar amino-acids found in at least 3 of the sequences. The secondary structural elements (helices as tubes, strands as arrows) of the yeast TmpK are also shown. Residues that contact TP_5A directly are marked with an arrow. The LID part of the sequence is marked as a dotted line; note the difference in the P-loop region between the eukaryotic TmpKs and the *E. coli* enzyme (between $\beta 1$ and $\alpha 1$), and the additional basic residues found only in *E. coli* TmpK.

Figure 5 Comparison of yeast thymidylate kinase (thick line) with HSV1 thymidine kinase (thin line) (pdb code 1kin). **a.** Superposition of both kinases (both are homodimers; only a monomer is shown) with TmpK in blue and HSV1-TK in yellow. The TP_5A bound to TmpK is also shown (cyan) as is the thymidine bound to HSV1-TK (orange). **b.** Stereo view (rotated 180° relative to the figure above) with a close-up of the active sites. Thymidine (orange) is displaced relative to the thymidine moiety of TP_5A (cyan). As a result, where in TmpK Tyr102 discriminates against ribonucleotides, the corresponding Tyr172 in

HSV1-TK makes a base-stacking interaction with the thymidine base, a role accomplished by Phe69 in TmpK.

Abbreviations:

TmpK, thymidylate kinase

UmpK, uridylate kinase

AK, adenylate kinase

HSV1-TK, herpes simplex virus thymidine kinase, type I

AZT, 3'-deoxy-3'-azido thymidine

AZT-MP, 3'-deoxy-3'-azido thymidine monophosphate

AZT-DP, 3'-deoxy-3'-azido thymidine diphosphate

AZT-TP, 3'-deoxy-3'-azido thymidine triphosphate

TP₅A, P1-(5'-adenosyl) P5-(5'-thymidyl) pentaphosphate

UP₅A, P1-(5'-adenosyl) P5-(5'-uridyl) pentaphosphate

PA, PB, PC, PD, PE, phosphate groups of TP₅A

NMP, nucleoside monophosphate kinases

ncs, non-crystallographic symmetry

RMSD, root mean square deviation

The examples illustrate the invention.

Thymidylate kinase (E.C. 2.7.4.9; ATP:dTMP phosphotransferase) catalyzes the phosphorylation of thymidine monophosphate (dTMP) to thymidine diphosphate (dTDP) utilizing ATP as its preferred phosphoryl donor (Jong & Campbell, 1984) according to the scheme:



Its location at the junction of the de-novo and salvage pathways for thymidine triphosphate (dTTP) synthesis makes thymidylate kinase (TmpK) an essential

enzyme for cell proliferation, and thus an attractive target for the development of drugs against cancer. In addition to its physiological role, TmpK is also involved in the activation of the AIDS drug 3'-azido-3'-deoxythymidine (AZT). AZT is a prodrug that must be phosphorylated three times to its triphosphate form (AZT-TP), since it is AZT-TP that inhibits viral replication by DNA chain termination. TmpK, which catalyzes the second phosphorylation step, from the monophosphate (AZT-MP) to the diphosphate (AZT-DP), has been shown to be the rate limiting enzyme in the AZT activation pathway (Furman et al., 1986, Qian et al., 1994). This results in a toxic accumulation of AZT-MP to millimolar concentration in cells exposed to AZT (Bridges et al., 1993, Tornevik et al., 1995, Yan et al., 1995), and in a low concentration of the active compound AZT-TP. Slow activation rates of prodrugs have been implicated in allowing the replicating virus to select for resistant mutants.

The Examples of the present invention provide the understanding the mechanism of phosphoryl transfer and pinpoint the amino acid residues involved and thus provide for a generally applicable method for developing and improving strategies for the treatment of cancer and AIDS. For example, the design of a mechanism-based inhibitor of TmpK resulting in the halt of dTTP synthesis, and thus cell proliferation, could play a role in chemotherapy of cancers.

Example 1: Structure of TmpK from yeast

TmpK from *Saccharomyces cerevisiae* (cdc8 gene), has 216 amino acid residues and a molecular weight of 25 kD. Crystals of the complex between the bisubstrate inhibitor TP5A with yeast thymidylate kinase were obtained by the hanging drop method. A 12 mg/mL enzyme solution was premixed with a TP5A solution to a final concentration of 10 mg/mL enzyme and 2 mM TP5A. Equal volumes of the protein-nucleotide solution and a solution composed of 20% PEG 2000 monomethyl ether, 100 mM sodium acetate pH 4.6, and 200 mM ammonium sulfate were mixed, and left to equilibrate at room temperature against a reservoir composed of the latter solution. Crystals with typical dimensions (μm) of $400 \times 200 \times 100$ grew within

days in space groups P1, P2₁, or P2₁2₁2 with very similar unit cell dimensions. This space group polymorphism made these crystals unsuitable for the initial structure determination (Lavie et al., 1997b). The structure reported here was solved by molecular replacement using the dTMP-TmpK complex structure as starting model (see below). A data set was collected at 100 K from a crystal soaked shortly in mother liquor with 25% glycerol as cryo-protectant, using a Siemens multiwire area detector mounted on a Mac Science rotating anode operating at 45kV, 100 mA. Processing of the data was carried out with XDS (Kabsch, 1993). The structure reported here is from a crystal which reduced in space group P2₁ and diffracted to 1.8 Å resolution, see Table 1.

Table 1: Data Collection and Refinement Statistics

Data collection

Temperature (K)	100
Resolution range (Å)	43.8 - 1.8
Observed reflections	270178
Unique reflections	116451
Completeness (% , overall/last shell)	82.0/56.9
R _{sym} ^a (% , overall/last shell)	5.9/26.6
Space group	P2 ₁
Unit-cell (Å, °)	a=72.6 b=87.3 c= 155.0 β=90.1
molecules / asymmetric unit	8

Refinement Statistics

Resolution range (Å)	43.8 - 2.0
Rfactor ^b /R _{free} (%)	20.9/27.9
RMS deviations	
bond lengths (Å)	0.012
bond angles (Å)	1.64
dihedral angles (°)	24.7
improper angles(°)	1.34
Reflections with F>0 s (working / test)	101931 / 5401

# of protein atoms	13791
nucleotide atoms	440
water molecules	1198
Average B (Å ²)	
main chain	20.5
side chain	22.1
waters	28.6
nucleotides	16.3
<hr/>	
a $R_{\text{sym}} = \frac{\sum_i I_i - \langle I \rangle }{\sum_i I_i}$	
b $R_{\text{factor}} = \frac{\sum_i F_{\text{obs}} - F_{\text{calc}} }{\sum_i F_{\text{obs}} }$ 5% of reflections were used for R_{free} .	
<hr/>	

The TP5A complex structure (Figure 1) was solved by the molecular replacement method with the partially refined dimeric dTMP complex structure as the starting search model. As mentioned above, the TP5A co-crystals exhibit space group polymorphism; the data set used has P2₁ symmetry with 4 dimers in the asymmetric unit. The pseudo-symmetry with the orthorhombic space group suggested the presence of two tetramers in the monoclinic asymmetric unit. This implied that when searching with the TmpK dimer, one should expect 2 rotation function solutions, where each solution has two different translation function solutions, thus yielding the two tetramers. Both AMoRe (Navaza, 1994) and X-PLOR (Brünger, 1993) yielded the two expected rotation functions solutions, but it was not possible to find the two appropriate translation function solutions. The self-Patterson map showed one clear peak that was interpreted as the vector relating the two tetramers. Thus, each of the two rotation function results were first applied on the search dimer, and then the translation vector from the self-Patterson was applied on the rotated dimers. The resulting two pairs of tetramers were used independently as search models for the translation function and yielded clear solutions. In P2₁ the origin of the unique axis is not defined so the relative position of one tetramer to the other was found by arbitrarily keeping one tetramer at y=0, which also limits the translational search to two dimensions (x and z). At first, the refinement was carried out using strict non-crystallographic symmetry (ncs) thus allowing to work with a dimer instead of the octamer. The refinement proceeded

smoothly with clear electron density for the TP5A bisubstrate inhibitor. When the R-factor reached 27% the ncs constraints were totally relaxed. The R-factor converged at 20% and 29% for the work and test set (5% of total reflections), respectively. Ncs restraints were reapplied at relatively low weight causing a slight decrease in R_{free} and a bigger increase in R_{work} . Consequently, the final model was refined with ncs-restraints grouping monomers 1, 3, 5, and 7 in one group and 2, 4, 6, and 8 in the other (the LID sequence and nucleotides were not included). The numbering used is residues 1 to 216 and 501 to 716 for monomer 1 and 2, respectively, with TP5A numbered 217 in monomer 1 and 717 in monomer 2. Monomers 3 and 4 are numbered as above with the addition of 1000, 5 and 6 with the addition of 2000, and 7 and 8 with the addition of 3000 (thus TP5A 3217 is the TP5A bound to monomer 7). Model building was done with the program O (Jones et al., 1991). The present model consists of 8 monomers with some residues omitted because of poor electron density (residues 1, 2, 1137 to 1148, 2001, 3001, 3501, 3502) and some residues modeled as alanines, 8 TP5A molecules, and 1198 water molecules (see Table 1). A simulated-annealing omit map (calculated without the TP5A and Arg15) with the current TP5A model superimposed is shown in Figure 2a.

Unlike the known structures of other nucleoside monophosphate (NMP) kinases (adenylate kinase, uridylate kinase, guanylate kinase) which are all monomeric, TmpK is a homodimer. Despite no amino acid sequence similarity between TmpK and any of the other NMP kinases, TmpK assumes the common fold of the other NMP kinases with 5 parallel beta strands forming a beta-sheet core surrounded by helices (Figure 1a). The highly hydrophobic dimer interface is composed of 3 parallel α -helices provided by each monomer that stack against each other.

The asymmetric unit of the P2₁ crystal form consists of eight monomers which pack in 4 dimers. A homodimer is the basic unit of yeast TmpK, but no cooperativity was detected in our kinetic assays (Lavie et al., 1997a). Upon overlaying all 8 monomers, it was evident that they can be divided into two groups; mon1, mon3, mon5 and mon7 adopt one conformation, whereas mon2, mon4,

mon6, and mon8 adopt another conformation (as neither ncs constraints nor restraints were used in the last stages of refinement, the conformation adopted by each monomer is totally independent of the other monomers; in the final cycle of refinement ncs-restraints were reapplied). For reasons that become apparent later, the first group was designated as inactive monomers, and the second group of monomers as active (Figure 1b). The RMSD between the active and inactive monomers excluding the LID sequence is 0.3 Å (on Ca atoms).

Most likely it is the crystal packing which determines which monomers adopt the active conformation, and which do not. Apparently, an active monomer of one dimer inactivates a monomer of another dimer. Arg173 (a non-conserved residue) from all active monomers interacts with the TP5A bound to the inactive monomers through a oxygen atom of phosphate PB, whereas Arg173 from the inactive monomers cannot form such an interaction. The presence of Arg173 prevents the formation of an active conformation of those monomers. Thus, it is this interaction of Arg173 from an active monomer with the TP5A of an inactive monomer which presumably prevents the closure of the LID of those monomers and results in poor density. Since TmpK is a dimer in solution (based on gel-filtration and dynamic light scattering experiments) and the fact that binding studies with TP5A clearly indicate a single class of binding sites in solution, it is assumed that the partitioning into active and inactive monomers is a pure crystallization artifact with no physiological significance. Therefore, the discussion is limited to the active conformers.

Example 2: Comparison with dTMP-TmpK complex structure

Kinases undergo large conformational changes upon the binding of substrates (Vornrhein et al., 1995). At least 3 different states have been described; an open state in the absence of substrates, a partially closed state when one of the substrates is bound, and a closed state when both substrates are present. The previously reported dTMP-TmpK complex (Lavie et al., 1997b) represents the partially closed state (ATP is missing) whereas the TP5A complex represents the

fully closed state, which is catalytically competent. Superposition of these two complexes shows that the main difference between them lies in the LID region (the other slight differences concentrate in loop sequences, which are expected to vary as a result of the different crystal packing between the two complexes). In NMP kinases the LID region has been observed to change its conformation upon ATP binding. In the dTMP-complex the LID sequence was only traceable in one of the monomers, and the density for the traced sequence was weak, indicating disorder, as expected in the absence of ATP. With TP5A it was expected to see a much more ordered LID region. In the monomers that were designated as the inactive conformer, the LID region is very hard to trace and lacks secondary structure. In contrast, the LID region of the active conformers was easier to trace and forms a helix (Figure 1a). Upon superposition of TmpK with uridylylate kinase (see more later for a detailed comparison with other NMP kinases), this newly formed helix overlays very well with a helix from the UmpK LID region (Scheffzek et al., 1996).

Example 3: Binding of TP₅A

Titration of TmpK to a cuvette containing 0.09 mM of the fluorescent bisubstrate inhibitor TP5A-MANT results in a fluorescence increase that indicates complex formation. Equilibrium fluorescence measurements were performed as described (Reinstein et al., 1990) using a SLM 8100 spectrofluorimeter with an excitation wavelength of 360 nm and emission wavelength of 440 nm. The experiments were done at pH 7.5 in a solution containing 50 mM Tris/HCl, 5 mM MgCl₂, 2 mM EDTA, and 100 mM KCl with the temperature of the cuvette being held constant at 25 °C. The fluorescent N-methylanthraniloyl group (MANT) joined to the ribose of the adenosine moiety of the bisubstrate inhibitor TP5A (TP5A-MANT) was used as probe (Reinstein et al., 1990).

Time resolved binding studies were performed with the same buffer as described above with a Hi-Tech Scientific SF61 stopped flow apparatus. As a signal for complex formation either the extrinsic signal of TP5A-MANT with excitation at 360 nm and a cutoff at 420 nm or the intrinsic tryptophan signal with excitation at 295

nm and a cutoff at 320 nm was used. These experiments were performed essentially as described (Packschies et al., 1997). The titration data were fitted by a quadratic equation (Reinstein et al., 1990) which yields a K_d of 135 nM for the binding of TP5A-MANT to TmpK. Competitive displacement of TP5A-MANT from the so formed complex by TP5A and analysis by a cubic equation (Thrall et al., 1996) shows the K_d for the binding of TP5A to TmpK to be 95 nM. The affinity of TP5A to TmpK appears to be rather weak compared to *E. coli* adenylate kinase that binds AP5A with a K_d of 15 nM or to *D. discoideum* UmpK that binds UP5A with a K_d of 3 nM (Reinstein et al., 1990, Wiesmüller et al., 1995). Time resolved measurements show that the binding of TP5A and TP5A-MANT is relatively fast but perhaps somewhat below the diffusion controlled limit with $k_{on} = 10 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$ and $k_{on} = 5.5 \times 10^6 \text{ M}^{-1} \text{ s}^{-1}$, respectively. This indicates that the conformational changes induced by the bisubstrate inhibitor, e.g. closing of the LID region over substrate, do not constitute the rate limiting step of the catalytic cycle since k_{cat} of yeast TmpK was shown to be 35 s^{-1} (Lavie et al., 1997a) whereas the binding of TP5A did not deviate from a linear relationship for k_{obs} versus [TP5A] up to 60 s^{-1} .

In the crystal the bisubstrate inhibitor TP5A appears to be bound tightly by all monomers (see Figure 2), as can be inferred from the relatively low B-factors (see Table 1) and the excellent electron density for the nucleotides. Before describing in detail the interactions made between the TP5A di-nucleotide and the enzyme, it is important to realize that the TP5A molecule is not a transition-state analog but is rather a bisubstrate or a biproduct analog (Scheffzek et al., 1996). In other words, it is not clear whether the observed TP5A is closer to the ATP-dTMP or the ADP-dTDP state.

The thymidine part of TP5A, which is completely surrounded by protein atoms, has a lower B-factor than the relatively exposed adenine part. The detailed interactions between the thymidine moiety and TmpK has been already described for the dTMP-TmpK complex (Lavie et al., 1997b), so only the differences in the TP5A complex will be outlined here. Lysine 37 is observed to interact with the phosphate of dTMP in the dTMP-TmpK complex structure. This phosphate (for phosphate

notation see figure 2c) corresponds to PE in TP5A. In the TP5A complex Lys37 interacts with PD instead; PD can be seen either as the connector between dTMP (PE) and ATP (PA, PB, PC), and thus as an artificial moiety, or as the b phosphate of dTDP. In the other NMP kinase structures with such bisubstrate analogs, the position of this connecting phosphate varies (Abele & Schulz, 1995, Müller & Schulz, 1992, Scheffzek et al., 1996). Thus, it is hard to interpret this change in Lys37 interaction from PE to PD, but it is possible that this residue participates in stabilizing the transferred phosphate group in the transition state.

The P-loop motif has been identified in many ATP- and GTP-binding proteins (Saraste et al., 1990). The binding of the nucleotide to the P-loop is through main-chain nitrogen atoms to the a and b phosphates of the nucleotide, and through a strictly conserved lysine. The ATP part of TP5A interacts in such a manner with the P-loop of TmpK, through the amides of Thr16 and Thr19 to the PB oxygen atoms, and Thr20 to PA oxygen atoms. In addition to the amide-phosphate interaction of Thr20, its side-chain also interacts with PA. The P-loop lysine (Lys18) is observed to interact with the PB and PC oxygen atoms of TP5A. It is positioned such that it may stabilize the transition state, in agreement with mutational analysis of adenylate kinase (Reinstein et al., 1988, Reinstein et al., 1990, Tsai & Yan, 1991). In the non-active monomers, Arg15 (situated at the tip of the P-loop) is in an extended conformation making no interaction with TP5A, while in the active monomers Arg15 bends to interact with a PC oxygen. It is in fact this interaction which lead the inventors dividing the monomers into active and inactive, since the inventors attribute an essential catalytic role to this arginine (see below).

The adenine part of TP5A makes only two interactions with the enzyme, both through the amino group at C6. One is to the main-chain carbonyl group of Lys187, the other through a water molecule to the highly conserved Gln21. This would explain the preference of TmpK for adenine nucleotides over guanine; in guanine-based nucleotides there is a carbonyl group at C6 instead of the amino group, which would prevent the favorable interaction with the main-chain carbonyl of Lys187. However, the partial acceptance of non-adenine-based nucleotides as phosphoryl donors by TmpK (Jong & Campbell, 1984) is consistent with our

structure, as there are no strong interactions between the adenine base and the enzyme, and there would be no steric discrimination against the amino group at the C2 position of guanine, which would face towards the solvent.

Bound magnesium was not observed in the structure, which is attributed to the low pH and the presence of ammonium sulfate in the crystallization conditions. Inferring from other NMP kinases (Scheffzek et al., 1996), the magnesium is expected to be octahedrally coordinated to the PB and PC oxygens (corresponding to b and g phosphates of ATP), Thr16, Asp93 (directly or through a water molecule), and two additional water molecules. At the position where magnesium is expected electron density was seen only for a water molecule bridging Asp93 and PC.

Example 4: Comparison with other NMP kinases

In addition to this TP5A-TmpK complex structure, there are at present 3 other structures of NMP kinases with bisubstrate analogs; adenylate kinase (yeast and *E.coli*) with AP5A (Abele & Schulz, 1995, Müller & Schulz, 1992), and uridylate kinase (slime mold) with UP5A (Scheffzek et al., 1996). The above mentioned adenylate kinases (AK) are both of the long LID region type, while uridylate kinase is more like TmpK having a short LID region (for the LID's amino acid sequence see Figure 4). Therefore, the inventors have chosen to compare TmpK to uridylate kinase (UmpK). The structure of UmpK complexed with UP5A was superimposed on the TP5A-TmpK complex structure by aligning the P-loop sequences of both enzymes. While UmpK and AK have a high degree of sequence identity, none exists between TmpK and either UmpK or AK, but structurally, TmpK overlays on UmpK surprisingly well and UP5A occupies nearly the same position as TP5A. There is, however, a striking difference. The LID domain of UmpK interacts directly with the phosphates of UP5A via the basic residues Arg131 and Arg137, which have been shown to stabilize the transition state (Schlichting & Reinstein, 1997). While lacking such basic residues in the LID domain, TmpK has Arg15 in the P-loop, the corresponding amino acid in UmpK being a glycine. And in fact, in what

the inventors defined as the monomers in the active conformation, Arg15 makes a 2.8 Å hydrogen-bond interaction with one oxygen atom of PC. This would be an example of the importance of having catalytic residues in proximity of the reaction center, but their origin, LID or P-loop, is irrelevant for catalysis.

Example 5: Comparison with Herpes Simplex Virus-1 thymidine kinase

From sequence similarity the Herpes Simplex Virus-1 thymidine kinase (HSV1-TK) has been grouped with the thymidylate kinase family. In fact, HSV1-TK has both a thymidine and a thymidylate kinase activity, suggesting that it evolved from thymidylate kinase and acquired the additional thymidine kinase activity. The 3-dimensional X-ray structure of HSV1-TK has been solved recently revealing, as for TmpK, a dimer as the basic unit (Brown et al., 1995). The viral kinase, however, contains 376 amino acids compared to TmpK's 216. It has additional N-terminal (45 residues) and C-terminal sequences (87 residues), as well as a few short inserts in the overlapping sequence region. Nevertheless, the overlapping sequence regions can be relatively easily overlaid structurally, with both kinases displaying the basic kinase fold as previously described (Figure 5).

The low level of substrate specificity of the viral kinase, which is medically important in the activation of nucleoside prodrugs, can now be rationalized upon comparison with the highly specific TmpK. Using the 14 amino-acid stretch of the P-loop regions of both enzymes for the superposition (Ca rms of 0.7 Å) it becomes apparent that the base moiety of thymidine bound to HSV1-TK is about 3.5 Å displaced in relation to the position of the base in TmpK. The displacement is away from the core of the enzyme towards the solvent. This slippage of the nucleotide towards the active site residues (mainly basic residues which would interact with phosphate groups) but without the concomitant contraction of the nucleotide's base binding pocket explains why HSV1-TK can phosphorylate even purines in addition to pyrimidines, and how it can possess both thymidine and thymidylate kinase activity. Presumably, after the first phosphorylation of thymidine to thymidine monophosphate, the nucleotide has room to slip deeper into the

binding site, thus placing the phosphate at the position previously occupied by the C5 hydroxyl. Now the second phosphorylation can take place. Thus, it is the nucleotide that moves relative to the enzyme's catalytic residues. This would suggest that HSV1-TK, while accepting purine nucleosides as substrates, might not accept purine monophosphates (the substrates for the thymidylate kinase activity) as they must be bound deeper into the binding cavity and thus might not fit. And in fact, the guanine based anti-herpes drug acyclovir is phosphorylated to the monophosphate by HSV1-TK, but the following phosphorylation step is catalyzed by guanylate kinase (Elion, 1982), consistent with this interpretation.

In contrast to the base, other residues in the active site overlay very well, noteworthy being Tyr102(TmpK) and Tyr172 (HSV1-TK) (within 0.3 Å). Tyr102 is responsible for the discrimination against ribonucleotides in TmpK by being located 3.5Å from the C2 of the ribose. In HSV1-TK it fulfills a completely different role by base stacking with thymine (in TmpK Phe69 stacks analogously to this Tyr with the base but from the opposite side of the base; see Figure 5b).

Important for the previous discussion of catalysis are the P-loop and LID regions; in this respect HSV1-TK is more similar to the *E.coli* TmpK than to the yeast TmpK in lacking the arginine in the P-loop (D14R15 in yeast, E12G13 and H58G59 in *E.coli* and HSV1-TK, respectively) and having basic residues in the LID region. Arg222 of HSV1-TK, which is in the LID region, is situated less than 4.5 Å from the deoxyribose O5' of thymidine. Analogously to other NMP kinases that utilize basic residues from the LID region to stabilize the negative charge developing in the transition state, we propose that Arg222 of HSV1-TK fulfills this catalytic role.

Example 6: Implications for catalysis and AZT activation

In addition to phosphorylating dTMP to dTDP, thymidylate kinase is also part of the activation pathway of the anti-HIV prodrug 3'-deoxy-3'-azidothymidine (AZT). In fact, TmpK is the rate-limiting enzyme in this activation pathway. Together with the previously reported nucleotide-complex structures with dTMP and with AZT-MP

(Lavie et al., 1997b), the TP5A-TmpK complex structure suggests which residues are needed for catalysis, and why AZT-MP is phosphorylated by TmpK so poorly. Whether the phosphoryl transfer mechanism is of an associative nature, a dissociative one, or of a mixed type, the negative charge developing during the transition state compared to the ground state has to be stabilized. This would be achieved by a basic residue (Arg or Lys), and its interaction with the phosphate must be made or strengthened at the transition state. In UmpK it has been shown that the mechanism of phosphoryl transfer is most probably associative, and that the charge developing on the transferred phosphoryl group is stabilized mainly by Arg131 and Arg137 from the LID region, in addition to Lys19, Arg148 and the catalytic magnesium ion. In contrast, yeast TmpK lacks basic residues in its LID region, but does have an arginine residue at the tip of the P-loop. In the TmpK-TP5A complex this arginine's side-chain is at a similar position as would be an arginine originating from the LID domain, and is located 2.8 Å from PC of TP5A. Therefore, the inventors attribute to Arg15 a similar role to that of the UmpK's LID arginines.

As pointed out previously (Lavie et al., 1997a, Lavie et al., 1997b), the binding of AZT-MP causes the P-loop of TmpK to shift by about 0.5 Å due to the interaction between the azido moiety of AZT and the side-chain of Asp14. This shift affects Arg15 as well, causing it to be not optimally located to fulfill its catalytic role. Most, but not all thymidylate kinases sequenced to date have an arginine in the P-loop. A noticeable exception is thymidylate kinase from *E. coli* which has a glycine residue at that position (like UmpK). The LID domain of the *E. coli* TmpK contains basic residues, unlike yeast TmpK but like UmpK. The present inventors postulate that the *E. coli* TmpK functions very similarly to the slime mold UmpK, where arginine residues from the LID domain participate in catalysis (candidates are Arg147, Arg149, Arg151, and Arg156). This would suggest that a P-loop movement of *E. coli* TmpK, like that observed with the yeast enzyme and AZT-MP, would not have such a detrimental effect on catalysis, as the residues important for catalysis (except the conserved lysine) do not originate from the P-loop. To test this

hypothesis, thymidylate kinase from *E. coli* was cloned, expressed and purified and its dTMP and AZT-MP phosphorylation rates were compared.

Example 7: Steady-state kinetics of *E. coli* TmpK

As predicted based on the structural rational explained above, the rate of phosphorylation of AZT-MP and dTMP by *E. coli* TmpK is comparable (Figure 3). In steady-state kinetics assays the catalytic activity of TmpK was measured with a coupled colorimetric assay essentially as described (Reinstein et al., 1988) with the following assay buffer: 100 mM Tris/HCl pH 7.5, 200 μ M NADH, 400 μ M phosphoenolpyruvate, 80 mM KCl, and 1.4 mM $MgCl_2$ at 25 °C. In addition, the assay contained different concentrations of ATP and dTMP and 10 to 50 nM TmpK as indicated in Figure 3. The data were analyzed with the Michaelis-Menten equation and the non-linear regression program Grafit (Erithacus software). While in the case of yeast TmpK AZT-MP is phosphorylated 200 fold slower than dTMP, for the *E. coli* enzyme the factor is only 2.5 (Table 2),

Table 2: Steady-state kinetic parameters

	yeast	<i>E. coli</i>
K_M for dTMP with ATP	9 μ M	2.7 μ M
K_M for AZTMP with ATP	6 μ M	30 μ M
K_M for ATP with dTMP	190 μ M	8 μ M
K_M for ATP with AZTMP	300 μ M	50 μ M
k_{cat} with ATP and dTMP	35 s ⁻¹	15 s ⁻¹
k_{cat} with ATP and AZTMP	0.175 s ⁻¹	6 s ⁻¹
ratio k_{cat} dTMP/AZTMP	200	2.5
ratio k_{cat}/K_M dTMP/AZTMP	133	27.5

illustrating the big difference in acceptance of AZT-MP by these two dTMP-kinases from different organisms. It appears from this comparison that NMP kinases have developed different strategies to attain catalysis; one class has catalytic residues in the P-Loop and the LID regions (AK, UmpK, etc.) whereas most dTMP kinases (*E. coli* TmpK being an exception) apparently lack the catalytic arginine residues of the LID region. In this respect it has also to be considered that the mechanism of phosphoryl transfer of the yeast TmpK may not be purely associative as was deduced for UmpK (Schlichting & Reinstein, 1997).

Example 8: Substitution of Arginine 15 with glycine in the P-loop of yeast TmpK and replacement of the yeast LID-region with the *E. coli* LID-region

In order to prove that the observations described in examples 1 to 7, supra, can be indeed applied to a nucleoside or nucleotide kinase naturally having only a low catalytic activity for nucleoside or nucleotide analogs, the yeast TmpK enzyme described above has been mutated at position X₃ in the consensus sequence of the P-loop (amino acid position 15 in the amino acid sequence of the yeast TmpK enzyme; see Figure 4) in order to substitute the amino acid arginine with glycine, the amino acid at the corresponding position in the P-loop of *E. coli* TmpK; see Figure 4. Furthermore, the LID sequence of the yeast TmpK enzyme from position 131 to 150 (FLSTQ ... GDER) was replaced by the corresponding *E. coli* TmpK LID sequence from position 138 to 150 (YLDVTP ... ELDR); see Figure 4. The mutations referred to above were introduced into the yeast TmpK either alone or in combination by introducing the corresponding modifications in the DNA sequence underlying the amino acid sequence of the yeast TmpK as described in Sambrook et al., supra. The resultant enzymes were then tested and compared with respect to dTMP and AZT-MP phosphorylation rates as described in Example 7 and in Reinstein et al., (1988) with 10 mM MgCl₂ and the following, constant concentration of nucleotides: 2 mM ATP and either 1 mM TMP or 1mM AZT-MP. Furthermore, as a control the TmpK enzymes from yeast, *E. coli*, human, mouse and herpes simplex virus were tested. The results are shown in Table 3 below.

Table 3

Protein	Activity TMP	Activity AZT-MP	Reduction
yeast	100	0.5	200
<i>E. coli</i>	43	21	2
yeast R → G	0.5	-	-
yeast → LID <i>E. coli</i>	4	0	-
yeast R → G + → LID <i>E. coli</i>	25	2	12
Human	5	0.07	70
Mouse	20	0.8	25
Herpes S.	1	0.5	2

As can be inferred from Table 3, the mutated version of the yeast TmpK enzyme that contains both, the mutated P-loop and LID-region exhibits improved kinase activity for the nucleotide analog AZT-MP which is 30-fold higher than that of the corresponding human enzyme. Moreover, the activity of the mutated yeast TmpK exceeds that from mouse and herpes simplex virus, the latter of which has been speculated to be useful in gene therapy.

Example 9: Improvement of kinase activity of human thymidylate kinase

Modifications at two different positions of a nucleoside monophosphate kinase sequence are listed as the means of achieving a kinase with enhanced prodrug phosphorylation activity. Those positions are either the P-loop region, the LID region, or both. The human thymidylate kinase (hTMPK) has been modified accordingly with the result that higher azidothymidine monophosphate (AZTMP) phosphorylation activity was achieved. In addition, a modification outside of the P-loop or the LID region, namely that of phenylalanine 105 to tyrosine, also results in a hTMPK variant having the desired kinetic activity with AZTMP. The mutations in the amino acid sequence of hTMPK (SEQ ID NO: 5) and testing of the resultant

polypeptides were performed as described in Example 8. The results are summarized in Table 4.

Table 4

Protein	Activity TMP	Activity AZTMP	ratio TMP/AZTMP
yeast TMPK (WT [†])	100 [‡]	0.5	200
A) human TMPK (WT)	1.9	0.019	100
B) Arg16Gly Small <i>E.coli</i> LID [§]	1.9	0.64	33
C) Arg16Gly Large <i>E. coli</i> LID [¶]	4.3	6.1	0.7
D) Phe105Tyr	0.475	0.665	0.7

[†]WT: wild-type

[‡]The rate of yeast TMPK with TMP is set at 100.

[§]Small *E. coli* LID: exchange of hTMPK LID sequence with that of *E. coli*, residues 145-148.

[¶]Large *E. coli* LID: exchange of hTMPK LID sequence with that of *E. coli*, residues 135-148.

The modifications of hTMPK in the P-loop (Arg16Gly) and in the LID region (large *E. coli* LID) have resulted in a variant (marked (C) in the table above) with 320 fold higher activity with AZTMP as substrate in comparison to the wild type hTMPK. Moreover, the substrate specificity has been changed dramatically, such that variant (C) phosphorylates AZTMP at a faster rate than the physiological substrate

TMP. This should improve its efficacy even further since TTP and AZTTP compete for incorporation into nascent viral DNA chains (e.g. TTP and AZTTP compete as substrates for reverse transcriptase). Variant (D), the result of a single amino-acid change, has likewise a faster rate with AZTMP than with TMP, but is about 10 times slower than variant (C).

The results obtained in accordance with the present invention are of far reaching medicinal importance since it for the first time enables a rational approach to using gene therapy to improve the antiviral impact of nucleoside and nucleotide analogs, such as AZT as outlined previously (Balzarini et al., 1988, Guettari et al., 1997, Lavie et al., 1997a).

References:

- Abele, Protein Sci 4 (1995), 1262-71.
- Balzarini, Biochem. Pharmacol. 37 (1988), 897-903.
- Bridges, Biochem Pharmacol 45 (1993), 1571-6.
- Brown, Nature Structural Biology 2 (1995), 876-81.
- Brünger, X-PLOR: a system for X-ray crystallography and NMR (1993), Yale University Press, New Haven, CT.
- Elion, American Journal of Medicine 73(1A) (1982), 7-13.
- Furman, Proc Natl Acad Sci U S A 83 (1986), 8333-7.
- Jones, Acta. Cryst. A 47 (1991), 110-119.
- Jong, J Biol Chem 259 (1984), 14394-8.
- Kabsch, J. Appl. Cryst. 24 (1993), 795-800.
- Lavie, Nature Medicine 3 (1997a), 922-924.
- Lavie, Nature Structural Biology 4 (1997b), 601-604.
- Müller, J Mol Biol 224 (1992), 159-77.
- Navaza, Acta Cryst. A 50 (1994), 157-163.
- Packschies, Biochemistry 36 (1997), 3417-22.
- Qian, Biochem Pharmacol 48 (1994), 383-9.
- Reinstein, Biochemistry 27 (1988), 4712-20.
- Reinstein, Biochemistry 29 (1990), 7440-50.
- Reynes, J Bacteriol 178 (1996), 2804-12.
- Saraste, Trends Biochem Sci 15 (1990), 430-4.
- Scheffzek Biochemistry 35 (1996), 9716-27.
- Schlichting, Biochemistry (1997).
- Studier Methods Enzymol 185 (1990), 60-89.
- Thrall, Biochemistry 35 (1996), 4609-18.
- Tornevik, Biochem Pharmacol 49 (1995), 829-37.
- Tsai, Biochemistry 30 (1991), 6806-18.
- Vonrhein, Structure 3 (1995), 483-90.
- Wiesmüller, Febs Lett 363 (1995), 22-4.
- Yan, J Biol Chem 270 (1995), 22836-41.

Abstract

Described are novel means and methods for the preparation and activation of nucleoside and nucleotide based drugs. In particular, a method for the production of a polypeptide having or having enhanced kinase activity for a nucleoside or nucleotide analog is provided and polypeptides obtainable by said method. Described are also polynucleotides and vectors encoding said polypeptide obtainable by the method of the invention as well as to host cells transformed therewith. Furthermore, antibodies against said polypeptide are provided and pharmaceutical and diagnostic compositions as well as kits comprising proteins having kinase activity for a nucleoside or nucleotide analog or their encoding polynucleotides or vectors. Furthermore, the use of the before described proteins, polypeptides, polynucleotides, vectors and antibodies is provided for the preparation of pharmaceutical compositions for treating, preventing and/or delaying a disease related to viral infection or cancer. In addition, a method for identifying inhibitors of nucleoside or nucleotide kinases is described and methods for identifying nucleoside or nucleotide based prodrugs employing the above mentioned polypeptides, polynucleotides, vectors and host cells. Also provided are compounds identifiable by said methods as well as pharmaceutical and diagnostic compositions comprising said inhibitors. Moreover, the use of proteins and polypeptides having nucleoside or nucleotide kinase activity or their encoding polynucleotides or vectors is described for the preparation of nucleoside or nucleotide phosphates or analogs and derivatives thereof.

ART 64/2001

17.04.00

PCT/EP99/00945


Max-Planck-Gesellschaft zur
Förderung der Wissenschaften e.V.
Our Ref.: B 3270 PCT

New set of claims

1. A method for the production of a polypeptide having or having enhanced kinase activity for a nucleoside or nucleotide analog, said method comprising substituting, adding or deleting at least one amino acid of a protein having nucleoside or nucleotide kinase activity at a position in the protein where:
 - (a) the amino acid is at position X_2 and/or X_3 in the consensus sequence $GX_1X_2X_3X_4GK$ of the P-loop;
 - (b) the amino acid is in the LID region; and/or
 - (c) the amino acid is at position 105 in the amino acid sequence of human thymidylate kinase or at a corresponding position in a protein having nucleoside or nucleotide kinase activity.
2. The method of claim 1, wherein said nucleoside is adenosine, cytidine, guanosine, thymidine or uridine or based on any of these.
3. The method of claim 1 or 2, wherein said nucleotide is a nucleoside monophosphate.
4. The method of claim 3, wherein said nucleoside monophosphate is thymidylate.
5. The method of any one of claims 1 to 4, wherein said protein is derived from a eukaryotic or prokaryotic organism.
6. The method of claim 5, wherein said organism is human or a yeast.
7. The method of any one of claims 1 to 6, wherein said protein comprises the amino acid sequence of any one of SEQ ID NOS: 1 to 13 or a fragment thereof.

4 17 04 00

8. The method of any one of claims 1 to 7, wherein said amino acid which is substituted or added in (a) is glutamic acid, glycine or lysine and the amino acid substituted in (c) is tyrosine.
9. The method of any one of claims 1 to 8, wherein said amino acid which is substituted or added in (b) is a basic amino acid, preferably arginine.
10. The method of any one of claims 1 to 9 where the LID has the consensus sequence R/KXXXXXERYEXXXQ.
11. The method of any one of claims 1 to 10, wherein said polypeptide exhibits kinase activity for a nucleoside or nucleotide analog which is higher than that of the/a corresponding wild type eukaryotic, preferably human enzyme.
12. The method of any one of claims 1 to 11, wherein said nucleoside analog is AZT, d4T or has the following structure:



wherein B is any nucleobase or analog thereof, and X is O, CH₂, NH or S.


13. The method of any one of claims 1 to 12, wherein the amino acid substitution(s) result in the P-loop and/or in the LID region of a bacterial nucleoside or nucleotide kinase, preferably those of the TmpK of E.coli.
14. A polynucleotide encoding the polypeptide obtainable by the method of any one of claims 1 to 13.
15. A vector containing the polynucleotide of claim 14.

17.04.00

16. The vector of claim 15, wherein the polynucleotide is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic cells.
17. The vector of claim 15 or 16, which is a gene transfer or a gene targeting vector.
18. A host cell genetically engineered with the vector of any one of claims 15 to 17.
19. A method for producing a polypeptide having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog comprising
- (a) culturing the host cell of claim 18, and
 - (b) recovering said polypeptide from the culture.
20. A method for producing cells capable of expressing a polypeptide having nucleoside or nucleotide kinase activity for nucleoside or nucleotide analogs comprising genetically engineering cells with the polynucleotide of claim 14, or with the vector of any one of claims 15 to 17.
21. A polypeptide having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog encoded by polynucleotide of claim 14, obtainable by the method of any one of claims 1 to 13 or 19 or from cells produced by the method of claim 20 or comprising a biologically active fragment of any of these.
22. An antibody specifically recognizing the polypeptide of claim 21.
23. A composition comprising
- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide; or



- (b) the polypeptide of claim 21, the polynucleotide of claim 14 or the vector of any one of claims 15 to 17;
 - (c) optionally a nucleoside or nucleotide analog; and
 - (d) optionally a pharmaceutically acceptable carrier.
24. The composition of claim 23, wherein said protein is a bacterial nucleoside or nucleotide kinase.
25. The composition of claim 23 or 24, wherein said protein is a bacterial TmpK.
26. The composition of any one of claims 23 to 25, wherein said protein has at least the P-loop and/or the LID region of E.coli TmpK.
27. The composition of claim 25 or 26, wherein said TmpK comprises the amino acid sequence shown in SEQ ID NO: 4 or a biologically active fragment thereof.
28. A kit comprising the polynucleotide of claim 14, the vector of any one of claims 15 to 17, the protein of claim 21 or the antibody of claim 22, and optionally a nucleoside or nucleotide analog.
29. A method for identifying an inhibitor of a nucleoside or nucleotide kinase comprising the steps of:
- (a) contacting the polypeptide of claim 21 or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a compound to be screened under conditions that permit binding of said compound to the nucleoside or nucleotide kinase, and
 - (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the absence or decrease of the signal is indicative for an inhibitor of a nucleoside or nucleotide kinase.
30. A method for identifying a nucleoside or nucleotide based prodrug comprising the steps of



- (a) contacting the polypeptide of claim 21 or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a nucleoside or nucleotide analog to be screened under conditions that permit kinase activity of said polypeptide, and
- (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the presence of a signal is indicative for a putative prodrug.
31. A method for the production of a pharmaceutical composition comprising the steps of
- (a) contacting the polypeptide of claim 21 or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a compound to be screened under conditions that permit binding of said compound to the nucleoside or nucleotide kinase, and
- (b) detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the absence or decrease of the signal is indicative for an inhibitor of a nucleoside or nucleotide kinase, or
- (a') contacting the polypeptide of claim 21 or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to kinase activity, with a nucleoside or nucleotide analog to be screened under conditions that permit kinase activity of said polypeptide, and
- (b') detecting presence or absence of a signal generated from the kinase activity of the polypeptide, wherein the presence of a signal is indicative for a putative prodrug; and
- (c) formulating the inhibitor identified in step (b) or the nucleoside or nucleotide analog identified in step (b') in a pharmaceutically acceptable form.

17.04.00

32. Use of
- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide,
 - (b) the polypeptide of claim 21, the polynucleotide of claim 14 or the vector of any one of claims 15 to 17; and/or
 - (c) the nucleoside or nucleotide analog identified in the method of claim 30 for the preparation of a pharmaceutical composition for the activation of nucleoside or nucleotide analogs or nucleoside or nucleotide based prodrugs and/or for the treatment of viral infections and/or diseases or cancer.
33. The use of claim 32, wherein said activation results in a cytotoxic nucleoside or nucleotide.
34. The use of claim 32 or 33, wherein said viral infection is HIV infection.
35. Use of the inhibitor obtainable by the method of claim 31 for the preparation of a pharmaceutical composition for inhibiting virus replication or for treating cancer.
36. A method for the preparative synthesis of a nucleoside phosphate analog comprising:
- (a) using a polynucleotide of claim 14 or as defined in any one of claim 23 to 27 in a noncellular system or in a cell *ex vivo*, ~~and~~
 - (b) formulating the cells modified in step (a) in a pharmaceutically acceptable form.
37. The composition of any one of claims 23 to 27 or the use of any one of claims 32 to 34, wherein said composition is a pharmaceutical composition and further comprises or is designed to be administered with a nucleoside or nucleotide analog, preferably AZT or d4T.

N 17.04.00

38. Use of

- (a) a prokaryotic protein having nucleoside or nucleotide kinase activity for a nucleoside or nucleotide analog or a polynucleotide encoding and capable of expressing said protein *in vivo* or a vector containing said polynucleotide,
- (b) the polypeptide of claim 21, the polynucleotide of claim 14 or the vector of any one of claims 15 to 17

for the preparation of nucleoside phosphates or analogs and derivatives thereof.

SEQUENCE LISTING

28. Juni 1999

<110> Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V.

<120> Novel means and methods for the preparation and
activation of nucleoside and nucleotide based drugs

<130> B3270PCT

<140> PCT/EP99/00945

<141> 1999-02-12

<150> EP 98 10 2546.3

<151> 1998-02-13

<160> 15

<170> PatentIn Ver. 2.1

<210> 1

<211> 240

<212> PRT

<213> African swine fever virus

<400> 1

Met Arg Gly Ile Leu Ile Thr Ile Glu Gly Ile Asn Gly Val Gly Lys
1 5 10 15

Ser Thr Gln Ala Met Arg Leu Lys Lys Ala Leu Glu Cys Met Asp Tyr
20 25 30

Asn Ala Val Cys Ile Arg Phe Pro Asn Pro Asp Thr Thr Thr Gly Gly
35 40 45

Leu Ile Leu Gln Val Leu Asn Lys Met Thr Glu Met Ser Ser Glu Gln
50 55 60

Leu His Lys Leu Phe Thr Lys His His Ser Glu Phe Ser Ala Glu Ile
65 70 75 80

Ala Ala Leu Leu Lys Leu Asn Phe Ile Val Ile Val Asp His Tyr Ile
85 90 95

Trp Ser Gly Leu Ala Tyr Ala Gln Ala Asp Gly Ile Thr Ile Glu Thr
100 105 110

Lys Asn Ile Phe Lys Pro Asp Tyr Thr Phe Phe Leu Ser Ser Lys Lys
115 120 125

Pro Leu Asn Glu Lys Pro Leu Thr Leu Gln Arg Leu Phe Glu Thr Lys
130 135 140

Glu Lys Gln Glu Thr Ile Phe Thr Asn Phe Thr Ile Ile Met Asn Asp
145 150 155 160

Val Pro Lys Asn Arg Leu Cys Ile Ile Pro Ala Thr Leu Asn Lys Glu
165 170 175

Ile Ile His Thr Met Ile Leu Thr Lys Thr Ile Lys Val Phe Asp Asn
180 185 190

Asn Ser Cys Leu Asn Tyr Ile Lys Met Tyr Asp Asp Lys Tyr Leu Asn
195 200 205

Val Gln Asp Leu Asn Leu Phe Asp Phe Asp Trp Gln Lys Cys Ile Glu
210 215 220

Asp Asn Asn Asp Lys Glu Glu Tyr Asp Asp Asp Asp Gly Phe Ile Ile
225 230 235 240

<210> 2

<211> 212

<212> PRT

<213> Bacillus subtilis

<400> 2

Met Ser Gly Leu Phe Ile Thr Phe Glu Gly Pro Glu Gly Ala Gly Lys
1 5 10 15

Thr Thr Val Leu Gln Glu Ile Lys Asn Ile Leu Thr Ala Glu Gly Leu
20 25 30

Gln Val Met Ala Thr Arg Glu Pro Gly Gly Ile Asp Ile Ala Glu Gln
35 40 45

Ile Arg Glu Val Ile Leu Asn Glu Asn Asn Ile Leu Met Asp Pro Lys
50 55 60

Thr Glu Ala Leu Leu Tyr Ala Ala Ala Arg Arg Gln His Leu Val Glu
65 70 75 80

Lys Val Lys Pro Ala Leu Glu Gln Gly Phe Ile Val Leu Cys Asp Arg
85 90 95

Phe Ile Asp Ser Pro Leu Ala Tyr Gln Gly Tyr Ala Arg Gly Leu Gly
100 105 110

Ile Asp Glu Val Leu Ser Ile Asn Glu Phe Ala Ile Gly Asp Met Met
115 120 125

Pro His Val Thr Val Tyr Phe Ser Ile Asp Pro Glu Glu Gly Leu Lys
130 135 140

Arg Ile Tyr Ala Asn Gly Ser Arg Glu Lys Asn Arg Leu Asp Leu Glu
145 150 155 160

Lys Leu Asp Phe His Thr Lys Val Gln Glu Gly Tyr Gln Glu Leu Met
165 170 175

Lys Arg Phe Pro Glu Arg Phe His Ser Val Asp Ala Gly Gln Ser Lys
180 185 190

Lys Glu Leu Asp Ala
210

<210> 4
 <211> 210
 <212> PRT
 <213> Haemophilus influenzae

<400> 4
 Met Lys Gly Lys Phe Ile Val Ile Glu Gly Leu Glu Gly Ala Gly Lys
 1 5 10 15
 Ser Ser Ala His Gln Ser Val Val Arg Val Leu His Glu Leu Gly Ile
 20 25 30
 Gln Asp Val Val Phe Thr Arg Glu Pro Gly Gly Thr Pro Leu Ala Glu
 35 40 45
 Lys Leu Arg His Leu Ile Lys His Glu Thr Glu Glu Pro Val Thr Asp
 50 55 60
 Lys Ala Glu Leu Leu Met Leu Tyr Ala Ala Arg Ile Gln Leu Val Glu
 65 70 75 80
 Asn Val Ile Lys Pro Ala Leu Met Gln Gly Lys Trp Val Val Gly Asp
 85 90 95
 Arg His Asp Met Ser Ser Gln Ala Tyr Gln Gly Gly Gly Arg Gln Leu
 100 105 110
 Asp Pro His Phe Met Leu Thr Leu Lys Glu Thr Val Leu Gly Asn Phe
 115 120 125
 Glu Pro Asp Leu Thr Ile Tyr Leu Asp Ile Asp Pro Ser Val Gly Leu
 130 135 140
 Ala Arg Ala Arg Gly Arg Gly Glu Leu Asp Arg Ile Glu Gln Met Asp
 145 150 155 160
 Leu Asp Phe Phe His Arg Thr Arg Ala Arg Tyr Leu Glu Leu Val Lys
 165 170 175
 Asp Asn Pro Lys Ala Val Val Ile Asn Ala Glu Gln Ser Ile Glu Leu
 180 185 190
 Val Gln Ala Asp Ile Glu Ser Ala Val Lys Asn Trp Trp Lys Ser Asn
 195 200 205
 Glu Lys
 210

<210> 5
 <211> 212
 <212> PRT
 <213> Homo sapiens

<400> 5
 Met Ala Ala Arg Arg Gly Ala Leu Ile Val Leu Glu Gly Val Asp Arg

1	5	10	15
Ala Gly Lys Ser Thr Gln Ser Arg Lys Leu Val Glu Ala Leu Cys Ala	20	25	30
Ala Gly His Arg Ala Glu Leu Leu Arg Phe Pro Glu Arg Ser Thr Glu	35	40	45
Ile Gly Lys Leu Leu Ser Ser Tyr Leu Gln Lys Lys Ser Asp Val Glu	50	55	60
Asp His Ser Val His Leu Leu Phe Ser Ala Asn Arg Trp Glu Gln Val	65	70	75
Pro Leu Ile Lys Glu Lys Leu Ser Gln Gly Val Thr Leu Val Val Asp	85	90	95
Arg Tyr Ala Phe Ser Gly Val Ala Phe Thr Gly Ala Lys Glu Asn Phe	100	105	110
Ser Leu Asp Trp Cys Lys Gln Pro Asp Val Gly Leu Pro Lys Pro Asp	115	120	125
Leu Val Leu Phe Leu Gln Leu Gln Leu Ala Asp Ala Ala Lys Arg Gly	130	135	140
Ala Phe Gly His Glu Arg Tyr Glu Asn Gly Ala Phe Gln Glu Arg Ala	145	150	155
Leu Arg Cys Phe His Gln Leu Met Lys Asp Thr Thr Leu Asn Trp Lys	165	170	175
Met Val Asp Ala Ser Lys Arg Leu Glu Ala Val His Glu Glu Leu Arg	180	185	190
Val Leu Ser Glu Asp Ala Ile Arg Thr Ala Thr Glu Lys Pro Leu Gly	195	200	205
Glu Leu Trp Lys	210		

<210> 6

<211> 188

<212> PRT

<213> Methanococcus jannaschii

<400> 6

Met Val Asp Asn Met Phe Ile Val Phe Glu Gly Ile Asp Gly Ser Gly	1	5	10	15
---	---	---	----	----

Lys Thr Thr Gln Ser Lys Leu Leu Ala Lys Lys Met Asp Ala Phe Trp	20	25	30
---	----	----	----

Thr Tyr Glu Pro Ser Asn Ser Leu Val Gly Lys Ile Ile Arg Glu Ile	35	40	45
---	----	----	----

Ser Leu Asp Trp Cys Lys Gln Pro Asp Val Gly Leu Pro Lys Pro Asp
115 120 125

Leu Ile Leu Phe Leu Gln Leu Gln Leu Leu Asp Ala Ala Ala Arg Gly
130 135 140

Glu Phe Gly Leu Glu Arg Tyr Glu Thr Gly Thr Phe Gln Lys Gln Val
145 150 155 160

Leu Leu Cys Phe Gln Gln Leu Met Glu Glu Lys Asn Leu Asn Trp Lys
165 170 175

Val Val Asp Ala Ser Lys Arg Thr Pro Ser Glu Thr Leu His Arg Gly
180 185 190

His Trp Gly Ser Tyr Gly Asn Lys Ser Ala Ser Ile Ala Asn Thr Ile
195 200 205

Phe Trp Phe Cys Lys Arg Leu Val Glu Gly Ser His Leu Tyr Thr Ile
210 215 220

Ser Arg Ser
225

<210> 8

<211> 210

<212> PRT

<213> Mycoplasma pneumoniae

<400> 8

Met Lys Gln Gly Val Phe Val Ala Ile Glu Gly Val Asp Gly Ala Gly
1 5 10 15

Lys Thr Val Leu Glu Ala Phe Lys Gln Arg Phe Pro Gln Ser Phe
20 25 30

Leu Gly Phe Lys Thr Leu Phe Ser Arg Glu Pro Gly Gly Thr Pro Leu
35 40 45

Ala Glu Lys Ile Arg Ala Leu Leu Leu His Glu Ala Met Glu Pro Leu
50 55 60

Thr Glu Ala Tyr Leu Phe Ala Ala Ser Arg Thr Glu His Val Arg Gln
65 70 75 80

Leu Ile Gln Pro Ala Leu Gln Gln Lys Gln Leu Val Ile Val Asp Arg
85 90 95

Phe Val Trp Ser Ser Tyr Ala Tyr Gln Gly Leu Ile Lys Lys Val Gly
100 105 110

Leu Asp Val Val Lys Lys Leu Asn Ala Asp Ala Val Gly Asp Ser Met
115 120 125

Pro Asp Phe Thr Phe Ile Val Asp Cys Asp Phe Glu Thr Ala Leu Asn
130 135 140

Arg Met Ala Lys Arg Gly Gln Asp Asn Leu Leu Asp Asn Thr Val Lys

145 150 155 160
 Lys Gln Ala Asp Phe Asn Thr Met Arg Gln Tyr Tyr His Ser Leu Val
 165 170 175
 Asp Asn Lys Arg Val Phe Leu Leu Asp Gly Gln Asn Gln Thr Gly Cys
 180 185 190
 Leu Glu Gln Phe Ile Glu Gln Leu Ser Gln Cys Leu Thr Gln Pro Thr
 195 200 205
 Leu Ser
 210

<210> 9
 <211> 210
 <212> PRT
 <213> Mycoplasma genitalium

<400> 9
 Met Asn Lys Gly Val Phe Val Val Ile Glu Gly Val Asp Gly Ala Gly
 1 5 10 15
 Lys Thr Ala Leu Ile Glu Gly Phe Lys Lys Leu Tyr Pro Thr Lys Phe
 20 25 30
 Leu Asn Tyr Gln Leu Thr Tyr Thr Arg Glu Pro Gly Gly Thr Leu Leu
 35 40 45
 Ala Glu Lys Ile Arg Gln Leu Leu Asn Glu Thr Met Glu Pro Leu
 50 55 60
 Thr Glu Ala Tyr Leu Phe Ala Ala Ala Arg Thr Glu His Ile Ser Lys
 65 70 75 80
 Leu Ile Lys Pro Ala Ile Glu Lys Glu Gln Leu Val Ile Ser Asp Arg
 85 90 95
 Phe Val Phe Ser Ser Phe Ala Tyr Gln Gly Leu Ser Lys Lys Ile Gly
 100 105 110
 Ile Asp Thr Val Lys Gln Ile Asn His His Ala Leu Arg Asn Met Met
 115 120 125
 Pro Asn Phe Thr Phe Ile Leu Asp Cys Asn Phe Lys Glu Ala Leu Gln
 130 135 140
 Arg Met Gln Lys Arg Gly Asn Asp Asn Leu Leu Asp Glu Phe Ile Lys
 145 150 155 160
 Gly Lys Asn Asp Phe Asp Thr Val Arg Ser Tyr Tyr Leu Ser Leu Val
 165 170 175
 Asp Lys Lys Asn Cys Phe Leu Ile Asn Gly Asp Asn Lys Gln Glu His
 180 185 190

Leu Glu Lys Phe Ile Glu Leu Leu Thr Arg Cys Leu Gln Gln Pro Thr
 195 200 205

His Tyr
 210

<210> 10

<211> 210

<212> PRT

<213> Schizosaccharomyces pombe

<400> 10

Met Ser Lys Gln Asn Arg Gly Arg Leu Ile Val Ile Glu Gly Leu Asp
 1 5 10 15

Arg Ser Gly Lys Ser Thr Gln Cys Gln Leu Leu Val Asp Lys Leu Ile
 20 25 30

Leu Asn Met Lys Arg Leu Lys Leu Phe Lys Phe Pro Asp Arg Thr Thr
 35 40 45

Ala Ile Gly Lys Lys Ile Asp Asp Tyr Leu Thr Glu Ser Val Gln Leu
 50 55 60

Asn Asp Gln Val Ile His Leu Leu Phe Ser Ala Asn Arg Trp Glu Pro
 65 70 75 80

Ser Ile Tyr Tyr Arg Ala Asn Gln Gln Arg Cys Asn Cys Ile Leu Asp
 85 90 95

Arg Tyr Ala Phe Ser Gly Ile Ala Phe Ser Ala Ala Lys Gly Leu Asp
 100 105 110

Trp Glu Trp Cys Lys Ser Pro Asp Arg Gly Leu Thr Arg Pro Asp Leu
 115 120 125

Val Ile Phe Leu Asn Val Asp Pro Arg Ile Ala Ala Thr Arg Gly Gln
 130 135 140

Tyr Gly Glu Glu Arg Tyr Glu Lys Ile Glu Met Gln Glu Lys Val Leu
 145 150 155 160

Lys Asn Leu Gln Arg Leu Gln Lys Glu Phe Arg Glu Glu Gly Leu Glu
 165 170 175

Phe Ile Thr Leu Asp Ala Ser Ser Tyr Ala Leu Glu Asp Val Asp Ser
 180 185 190

Gln Ile Val Asp Leu Val Ser Asn Val Asn Ile His Glu Thr Leu Asp
 195 200 205

Val Leu
 210

<210> 11

<211> 204
 <212> PRT
 <213> Vaccinia virus

<400> 11

Met Ser Arg Gly Ala Leu Ile Val Phe Glu Gly Leu Asp Lys Ser Gly
 1 5 10 15

Lys Thr Thr Gln Cys Met Asn Ile Met Glu Ser Ile Pro Ala Asn Thr
 20 25 30

Ile Lys Tyr Leu Asn Phe Pro Gln Arg Ser Thr Val Thr Gly Lys Met
 35 40 45

Ile Asp Asp Tyr Leu Thr Arg Lys Lys Thr Tyr Asn Asp His Ile Val
 50 55 60

Asn Leu Leu Phe Cys Ala Asn Arg Trp Glu Phe Ala Ser Phe Ile Gln
 65 70 75 80

Glu Gln Leu Glu Gln Gly Ile Thr Leu Ile Val Asp Arg Tyr Ala Phe
 85 90 95

Ser Gly Val Ala Tyr Ala Ala Ala Lys Gly Ala Ser Met Thr Leu Ser
 100 105 110

Lys Ser Tyr Glu Ser Gly Leu Pro Lys Pro Asp Leu Val Ile Phe Leu
 115 120 125

Glu Ser Gly Ser Lys Glu Ile Asn Arg Asn Val Gly Glu Glu Ile Tyr
 130 135 140

Glu Asp Val Thr Phe Gln Gln Lys Val Leu Gln Glu Tyr Lys Lys Met
 145 150 155 160

Ile Glu Glu Gly Asp Ile His Trp Gln Ile Ile Ser Ser Glu Phe Glu
 165 170 175

Glu Asp Val Lys Lys Glu Leu Ile Lys Asn Ile Val Ile Glu Ala Ile
 180 185 190

His Thr Val Thr Gly Pro Val Gly Gln Leu Trp Met
 195 200

<210> 12
 <211> 205
 <212> PRT
 <213> Variola virus

<400> 12

Met Ser Arg Gly Ala Leu Ile Val Phe Glu Gly Leu Asp Lys Ser Gly
 1 5 10 15

Lys Thr Thr Gln Cys Met Asn Ile Met Glu Ser Ile Pro Thr Asn Thr
 20 25 30

Ile Lys Tyr Leu Asn Phe Pro Gln Arg Ser Thr Val Thr Gly Lys Met
35 40 45

Ile Asp Asp Tyr Leu Thr Arg Lys Lys Thr Tyr Asn Asp His Ile Val
50 55 60

Asn Leu Leu Phe Cys Ala Asn Arg Trp Glu Phe Ala Ser Phe Ile Gln
65 70 75 80

Glu Gln Leu Glu Gln Gly Ile Thr Leu Ile Val Asp Arg Tyr Ala Phe
85 90 95

Ser Gly Val Ala Tyr Ala Thr Ala Lys Gly Ala Ser Met Thr Leu Ser
100 105 110

Lys Ser Tyr Glu Ser Gly Leu Pro Lys Pro Asp Leu Val Ile Phe Leu
115 120 125

Glu Ser Gly Ser Lys Glu Ile Asn Arg Asn Val Gly Glu Glu Ile Tyr
130 135 140

Glu Asp Val Ala Phe Gln Gln Lys Val Leu Gln Glu Tyr Lys Lys Met
145 150 155 160

Ile Glu Glu Gly Glu Asp Ile His Trp Gln Ile Ile Ser Ser Glu Phe
165 170 175

Glu Glu Asp Val Lys Lys Glu Leu Ile Lys Asn Ile Val Ile Glu Ala
180 185 190

Ile His Thr Val Thr Gly Pro Val Gly Gln Leu Trp Met
195 200 205

<210> 13

<211> 216

<212> PRT

<213> *Saccharomyces cerevisiae*

<400> 13

Met Met Gly Arg Gly Lys Leu Ile Leu Ile Glu Gly Leu Asp Arg Thr
1 5 10 15

Gly Lys Thr Thr Gln Cys Asn Ile Leu Tyr Lys Lys Leu Gln Pro Asn
20 25 30

Cys Lys Leu Leu Lys Phe Pro Glu Arg Ser Thr Arg Ile Gly Gly Leu
35 40 45

Ile Asn Glu Tyr Leu Thr Asp Ser Phe Gln Leu Ser Asp Gln Ala
50 55 60

Ile His Leu Leu Phe Ser Ala Asn Arg Trp Glu Ile Val Asp Lys Ile
65 70 75 80

Lys Lys Asp Leu Leu Glu Gly Lys Asn Ile Val Met Asp Arg Tyr Val
85 90 95

Tyr Ser Gly Val Ala Tyr Ser Ala Ala Lys Gly Thr Asn Gly Met Asp
 100 105 110

Leu Asp Trp Cys Leu Gln Pro Asp Val Gly Leu Leu Lys Pro Asp Leu
 115 120 125

Thr Leu Phe Leu Ser Thr Gln Asp Val Asp Asn Asn Ala Glu Lys Ser
 130 135 140

Gly Phe Gly Asp Glu Arg Tyr Glu Thr Val Lys Phe Gln Glu Lys Val
 145 150 155 160

Lys Gln Thr Phe Met Lys Leu Leu Asp Lys Glu Ile Arg Lys Gly Asp
 165 170 175

Glu Ser Ile Thr Ile Val Asp Val Thr Asn Lys Gly Ile Gln Glu Val
 180 185 190

Glu Ala Leu Ile Trp Gln Ile Val Glu Pro Val Leu Ser Thr His Ile
 195 200 205

Asp His Asp Lys Phe Ser Phe Phe
 210 215

<210> 14

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic, no
 natural origin, hypothetical

<400> 14

ggaattccat atgcgcagta agtatatcgt c

31

<210> 15

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic, no
 natural origin, hypothetical

<400> 15

cgcggtatcct catgcgtcca actcattcac ccag

34




Fig. 1




a**b**

Figure 2






Fig. 3 (cont.)




Fig. 4




Figure 4




Figure 4 (Cont.)



a**b**

FIGURE 5

DECLARATION
Utility Application



Lyon & Lyon LLP
Docket Information
256/137

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **NOVEL MEANS AND METHODS FOR THE PREPARATION AND ACTIVATION OF NUCLEOSIDE AND NUCLEOTIDE BASED DRUGS** the specification of which

(Check One)

☐
☒

is attached hereto OR
was filed on 08/10/00/ as United States Application Serial No. 09/622,101
or PCT International Application No. PCT/EP00/0945 and was amended on
April 17, 2000 (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Date of Filing	Priority Claimed	
			Yes	No
EP 98 10 2546.3	DE, et al.	Feb. 13, 1998	Yes	

I hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date	Status-Patented, Pending or Abandoned
	PCT/EP99/0945	Feb. 12, 1999	

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Title 18, United States Code, § 1001 and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

201	FULL NAME OF INVENTOR	FIRST Name <u>Roger</u>	MIDDLE Initial <u>S.</u>	LAST Name <u>GOODY</u>	
	RESIDENCE & CITIZENSHIP	City <u>Dortmund</u>	State or Foreign Country <u>GERMANY</u>	Country of Citizenship <u>GERMANY</u>	
	POST OFFICE ADDRESS	<u>Harnackstrasse 61b</u>	City <u>Dortmund</u>	State or Country <u>GERMANY</u>	Zip Code <u>44139</u>
INVENTOR'S SIGNATURE <u>X [Signature]</u>					DATE <u>02. Nov. 2000</u>

202	FULL NAME OF INVENTOR	FIRST Name <u>Manfred</u>	MIDDLE Initial	LAST Name <u>KONRAD</u>	
	RESIDENCE & CITIZENSHIP	City <u>Göttingen</u>	State or Foreign Country <u>GERMANY</u>	Country of Citizenship <u>GERMANY</u>	
	POST OFFICE ADDRESS	<u>Zur Scharfmühle 74</u>	City <u>Göttingen</u>	State or Country <u>GERMANY</u>	Zip Code <u>37083</u>
INVENTOR'S SIGNATURE <u>[Signature]</u>					DATE <u>09. Nov. 2000</u>

203	FULL NAME OF INVENTOR	FIRST Name <u>Arnon</u>	MIDDLE Initial	LAST Name <u>LAVIE</u>	
	RESIDENCE & CITIZENSHIP	City <u>Chicago</u>	State or Foreign Country <u>il</u>	Country of Citizenship <u>GERMANY</u>	
	POST OFFICE ADDRESS	<u>College of Medicine, Dept. of Biochemistry and Molecular Biology (MC536), 1819 West Polk Street</u>	City <u>Chicago</u>	State or Country <u>IL</u>	Zip Code <u>60612</u>
INVENTOR'S SIGNATURE <u>A. Lavie</u>					DATE <u>15. Nov. 2000</u>

204	FULL NAME OF INVENTOR	FIRST Name <u>Joachim</u>	MIDDLE Initial	LAST Name <u>REINSTEIN</u>	
	RESIDENCE & CITIZENSHIP	City <u>Dortmund</u>	State or Foreign Country <u>GERMANY</u>	Country of Citizenship <u>GERMANY</u>	
	POST OFFICE ADDRESS	<u>Plauener Strasse 54</u>	City <u>Dortmund</u>	State or Country <u>GERMANY</u>	Zip Code <u>44139</u>
INVENTOR'S SIGNATURE <u>X [Signature]</u>					DATE <u>1 Nov. 2000</u>

205	FULL NAME OF INVENTOR	FIRST Name <u>Ilme</u>	MIDDLE Initial <u>S</u>	LAST Name <u>SCHLICHTING</u>	
	RESIDENCE & CITIZENSHIP	City <u>Dortmund</u>	State or Foreign Country <u>GERMANY</u>		Country of Citizenship <u>GERMANY</u>
	POST OFFICE ADDRESS	Redtenbacher Strasse 30	City Dortmund	State or Country GERMANY	Zip Code 44139
INVENTOR'S SIGNATURE <u>X Ilme Schlichting</u>			DATE <u>2 Nov 2000</u>		